

# THE MODULI STACK OF STABLE RELATIVE IDEAL SHEAVES

BAOSEN WU

**ABSTRACT.** In this paper, we propose a definition of the moduli stack of stable relative ideal sheaves, and prove that it is a separated and proper Deligne-Mumford stack. It is the first part of the project of relative Donaldson-Thomas theory of ideal sheaves on projective threefolds, which in the end provides a degeneration formula of Donaldson-Thomas invariants of threefolds.

## INTRODUCTION

Donaldson-Thomas invariants are defined via integrals over the virtual fundamental class of the moduli space of stable sheaves [3, 12]. Recently, D. Maulik, N. Nekrasov, A. Okounkov and R. Pandharipande [10, 11] investigated the moduli scheme of ideal sheaves of one dimensional closed subschemes in smooth projective threefolds and proposed conjectures relating Donaldson-Thomas invariants with Gromov-Witten invariants. They also verified these conjectures in the case of local toric Calabi-Yau threefolds.

Despite the efforts made by many people, the calculation of Donaldson-Thomas invariants for general threefolds is still difficult. The current main tool for such calculation is the localization formula of virtual fundamental class, which is originally developed for the calculation of Gromov-Witten invariants. Another important tool is the technique of degeneration, which has been applied to Gromov-Witten theory successfully. The algebraic geometric machinery of degeneration in moduli problems has been developed by Jun Li [7] in order to prove a degeneration formula for Gromov-Witten invariants. It is believed that there is a similar degeneration formula in Donaldson-Thomas theory.

The goal of this paper is to define the moduli stack of stable relative ideal sheaves which lays the foundation of relative Donaldson-Thomas theory. In a subsequent paper, by applying the theory of virtual cycles, Jun Li and the author will prove a degeneration formula relating Donaldson-Thomas invariants of a smooth threefold with relative Donaldson-Thomas invariants of pairs.

Now we recall briefly the definition of Donaldson-Thomas invariants of Calabi-Yau threefolds via moduli scheme of ideal sheaves. Let  $X$  be a smooth projective Calabi-Yau threefold with  $\beta \in H_2(X, \mathbf{Z})$ . There is a projective scheme  $I_n(X, \beta)$  parameterizing ideal sheaves  $\mathcal{I}_Z$  of one dimensional closed subscheme  $Z \subset X$  with  $[Z] = \beta$  and  $\chi(\mathcal{O}_Z) = n$ . Following the theory of virtual fundamental cycles [1, 6], R. P. Thomas [12] showed that there exists a virtual fundamental class in the moduli scheme of stable sheaves on a Calabi-Yau threefold. Applying this result to the moduli scheme  $I_n(X, \beta)$ , he obtained a virtual fundamental class  $[I_n(X, \beta)]^{vir}$ . By Serre duality, the virtual dimension of  $I_n(X, \beta)$  is 0. Therefore, one can define the Donaldson-Thomas invariant  $N_n(\beta) = \deg[I_n(X, \beta)]^{vir}$ .

Next we explain briefly our approach in this paper. To begin with, we recall the setup of the machinery of degeneration in moduli problems. Let  $C$  be a smooth affine curve which serves as a parameter space. Fix a closed point  $\mathbf{0} \in C$ . Let  $\pi : X \rightarrow C$  be a flat family of projective schemes such that the total space  $X$  is smooth and the fibers  $X_t$  are smooth varieties for  $t \neq 0$ , and  $X_0$  is a union of two smooth irreducible varieties  $Y_1$  and  $Y_2$  intersecting transversally along a smooth divisor  $D$ . Since the original Donaldson-Thomas invariant is only defined for smooth projective threefolds. To apply this degeneration technique, the immediate goal is to construct a “good” moduli of ideal sheaves on  $X_0$  so that its obstruction theory is perfect, and the associated Donaldson-Thomas invariant of  $X_0$  is well defined which is equal to that of  $X_t$ .

Naturally, one can consider the moduli space of rank 1 stable sheaves on  $X_0$ . To be precise, fix a relative ample line bundle  $H$  over  $X/C$ . As is well known, there is a projective scheme parameterizing rank 1 stable sheaves on  $X/C$ . On a smooth fibre  $X_t$ , a rank 1 stable sheaf is essentially an ideal sheaf and hence Donaldson-Thomas invariants are well defined. For the singular fibre  $X_0$ , although the moduli scheme is well defined, we still need the existence of virtual fundamental classes. Unfortunately, the standard tangent obstruction theory of this moduli problem is in general not perfect and hence we don’t know the existence of the virtual cycle.

To overcome this difficulty, we borrow the notion of stack of expanded degeneration  $\mathfrak{X}$  introduced by Jun Li [7] to replace the scheme  $X/C$ , and construct the moduli stack  $\mathfrak{I}_{X/C}^\Gamma$  of stable perfect ideal sheaves on  $\mathfrak{X}$ . The price we pay is to work on stacks rather than on schemes. Now we explain the idea in a little more detail. The difference between the family  $X/C$  and the stack  $\mathfrak{X}$  appears only in the central fiber  $X_0$  and  $\mathfrak{X}_0$ , where  $X_0$  is a union of two smooth irreducible varieties, while  $\mathfrak{X}_0$  is a groupoid of all semistable models of  $X_0$ . Here a semistable model  $X[n]_0$  is a chain of smooth varieties  $Y_1, \Delta_1, \dots, \Delta_n, Y_2$  intersecting transversally along smooth divisors which are all isomorphic to  $D$ , and  $\Delta_i \cong \Delta$  is a projective bundle  $\mathbf{P}(N \oplus \mathbf{1})$  over  $D$ , where  $N$  is the normal line bundle of  $D$  in  $Y_2$  and  $\mathbf{1}$  is the trivial line bundle over  $D$ . Instead of rank 1 stable sheaves on  $X_0$ , we consider *perfect* ideal sheaves on semistable models of  $X_0$ . The rigorous definition of perfect ideal sheaves is given in section 2. When the relative dimension of  $X/C$  is not greater than three, a perfect ideal sheaf on  $X_0$  has vanishing local obstruction to deforming to ideal sheaves on smooth nearby fibres of  $X_0$ . By using stable perfect ideal sheaves, we get a perfect tangent obstruction theory on this moduli problem. However, this moduli scheme of perfect ideal sheaves on  $X_0$  is no longer proper. To compactify the moduli space, we include stable perfect ideal sheaves on semistable models of  $X_0$  as well. This compactified moduli  $\mathfrak{I}_{X/C}^\Gamma(\mathbf{0})$  is in fact a Deligne-Mumford stack rather than a scheme.

As a preparation for the degeneration formula, we need to decompose the moduli stack  $\mathfrak{I}_{X/C}^\Gamma(\mathbf{0})$  according to the decomposition of  $X_0$  into two pairs  $(Y_i, D_i)$ . In general, a semistable model of  $X_0$  can be decomposed into two parts, which are called extended pairs. As a result, a perfect ideal sheaf on  $X[n]_0$  can be regarded as the gluing of two ideal sheaves on the extended pairs which are called relative ideal sheaves. Indeed,  $\mathfrak{I}_{X/C}^\Gamma(\mathbf{0})$  is canonically a finite union of fibre products of moduli stacks  $\mathfrak{I}_{Y_1, D_1}^{\Gamma_1}$  and  $\mathfrak{I}_{Y_2, D_2}^{\Gamma_2}$  of stable relative ideal sheaves.

The main results of this paper are the following theorems.

**Theorem 0.1.** *The moduli stack  $\mathfrak{I}_{X/C}^\Gamma$  of stable perfect ideal sheaves of topological type  $\Gamma \in H_*(X, \mathbf{Z})$  on  $\mathfrak{X}$  is a separated and proper Deligne-Mumford stack of finite type over  $C$ .*

**Theorem 0.2.** *The moduli stack  $\mathfrak{I}_{Y,D}^\Gamma$  of stable relative ideal sheaves of topological type  $\Gamma \in H_*(Y, \mathbf{Z})$  on  $\mathfrak{Y}$  is a separated and proper Deligne-Mumford stack of finite type.*

We take a look at one interesting application of the degeneration formula of Donaldson-Thomas invariants in the case  $\beta = 0$ . For any smooth projective three-fold  $X$ , the moduli space  $I_n(X, 0)$  always has virtual dimension 0 and hence  $N_n(0)$  is well defined. A conjecture [11] says that

$$\sum_{k=0}^{\infty} N_k(0) q^k = M(-q)^{\int_X c_3(T_X \otimes K_X)},$$

where  $M(q) = \prod_{n \geq 1} \frac{1}{(1-q^n)^n}$  is the McMahon function. This conjecture has been proved by Jun Li in [9]. He showed that the zero dimensional Donaldson-Thomas invariants depend only on the Chern numbers of  $X$  and by complex cobordism theory, the problem can be reduced to toric cases  $X = \mathbf{P}^3, \mathbf{P}^2 \times \mathbf{P}^1, \mathbf{P}^1 \times \mathbf{P}^1 \times \mathbf{P}^1$  which are all known. Later, M. Levine and R. Pandharipande [5] gave a different argument based on algebraic cobordism theory and the assumption of the degeneration formula of zero dimensional Donaldson-Thomas invariants.

This paper is organized as follows. In section 1 we review the notion of expanded degenerations of  $X/C$ . For a detailed description and proof of the results, the reader is encouraged to read the original paper [7]. In section 2, we define perfect ideal sheaves and the stability condition, and prove the basic properties of the stack of stable perfect ideal sheaves. The proof of the properness of this stack relies on a key technical lemma which is the main content of section 4. In section 3, we study the moduli stack of stable relative ideal sheaves, and show that similar results hold as in section 2. Finally we state the relation between the moduli stack of stable perfect ideal sheaves on  $\mathfrak{X}_0$  and the moduli stack of stable relative ideal sheaves on pairs  $\mathfrak{Y}_i$ .

I am most grateful to my advisor Jun Li for sharing the idea with me. His help for the preparation of this paper is invaluable.

## 1. THE STACK OF EXPANDED DEGENERATION

Let  $C$  be a nonsingular affine curve with a distinguished closed point  $\mathbf{0} \in C$ . Let  $\pi : X \rightarrow C$  be a flat projective family of schemes of relative dimension  $d > 0$ , which satisfies the following conditions:

- (1)  $X$  is smooth and  $\pi$  is smooth away from the central fiber  $X_0 = X \times_C \mathbf{0}$ ;
- (2) The scheme  $X_0$  is a union of two smooth irreducible components  $Y_1$  and  $Y_2$  intersecting transversally along a smooth divisor  $D$ .

In this section, we will review the definition and some properties of the Artin stack  $\mathfrak{X}$  of expanded degeneration of  $X/C$ . Now we begin with standard models.

**1.1. The standard model.** Let  $G_m$  be the multiplicative group of the ground field  $k = \mathbf{C}$ . Let  $G[n]$  denote the algebraic group  $G_m \times \cdots \times G_m$  ( $n$  copies). We will define schemes  $C[n]$  and  $X[n]$  with  $G[n]$  actions, and a  $G[n]$ -equivariant morphism  $\pi_n : X[n] \rightarrow C[n]$ .

First, we describe the base scheme  $C[n]$  and a  $G[n]$  action on  $C[n]$ . We fix isomorphisms  $\mathbf{A}^{n+1} \cong \operatorname{Spec} k[t_1, \dots, t_{n+1}]$ ,  $\mathbf{A}^1 \cong \operatorname{Spec} k[s]$  and  $G_m \cong \operatorname{Spec} k[\sigma, \sigma^{-1}]$ . These isomorphisms give distinguished closed points  $\mathbf{0} \in \mathbf{A}^{n+1}$  and  $\mathbf{0} \in \mathbf{A}^1$ . There is a  $G[n]$  action on  $\mathbf{A}^{n+1}$  defined as follows,

$$(t_1, \dots, t_{n+1})^\sigma = (\sigma_1 t_1, \sigma_1^{-1} \sigma_2 t_2, \dots, \sigma_{n-1}^{-1} \sigma_n t_n, \sigma_n^{-1} t_{n+1}),$$

where  $\sigma = (\sigma_1, \dots, \sigma_n) \in G[n]$  and  $(t_1, \dots, t_{n+1}) \in \mathbf{A}^{n+1}$ . Let  $\mathbf{p} : \mathbf{A}^{n+1} \rightarrow \mathbf{A}^1$  be a morphism defined as

$$\mathbf{p}(t_1, \dots, t_{n+1}) = t_1 t_2 \cdots t_{n+1}.$$

Then  $\mathbf{p}$  is  $G[n]$ -equivariant with the trivial  $G[n]$ -action on  $\mathbf{A}^1$ . Let  $A^* = \mathbf{A}^1 - \{\mathbf{0}\}$ . Then  $\mathbf{p}^{-1}(A^*)$  is a principal  $G[n]$ -bundle over  $A^*$ .

Let  $C$  be the nonsingular affine curve. Fix once and for all an étale morphism  $C \rightarrow \mathbf{A}^1$  such that the inverse image of  $\mathbf{0} \in \mathbf{A}^1$  is a set of the single distinguished point  $\mathbf{0} \in C$ . Let  $C[n] = C \times_{\mathbf{A}^1} \mathbf{A}^{n+1}$ . Then  $C[n]$  has a distinguished closed point lying over the distinguished points of  $C$  and  $\mathbf{A}^{n+1}$ , and there is a well-defined  $G[n]$  action on  $C[n]$  induced from the  $G[n]$  action on  $\mathbf{A}^{n+1}$  and the trivial action on  $C$ . Let  $p : C[n] \rightarrow C$  be the projection to the first factor. Let  $C^* = C - \{\mathbf{0}\}$  and  $C[n]^* = p^{-1}(C^*)$ . Then  $p : C[n]^* \rightarrow C^*$  is a principal  $G[n]$ -bundle over  $C^*$ .

Let  $[n+1] = \{1, 2, \dots, n+1\}$  be a set of integers. Let  $m \leq n$  be a nonnegative integer and  $I \subset [n+1]$  be a subset with cardinality  $|I| = m+1$ . There is a unique strictly increasing function  $\iota_I : [m+1] \rightarrow [n+1]$  whose image in  $[n+1]$  is  $I$ . Fix an isomorphism  $\mathbf{A}^{m+1} \cong \operatorname{Spec} k[w_1, \dots, w_{m+1}]$ . A standard embedding  $\gamma_I : \mathbf{A}^{m+1} \rightarrow \mathbf{A}^{n+1}$  is defined by a homomorphism  $k[t_1, \dots, t_{n+1}] \rightarrow k[w_1, \dots, w_{m+1}]$  as

$$t_l \mapsto \begin{cases} w_k & \text{if } l = \iota_I(k) \in I \\ 1 & \text{if } l \notin I. \end{cases}$$

Since this embedding is compatible with the projection  $\mathbf{p} : \mathbf{A}^{n+1} \rightarrow \mathbf{A}^1$ , it induces a standard embedding

$$\gamma_I : C[m] \longrightarrow C[n].$$

There is another embedding  $\mathbf{c}_I : C[m] \rightarrow C[n]$  corresponding to a coordinate plane of  $C[n]$ , which is defined by

$$t_l \mapsto \begin{cases} w_k & \text{if } l = \iota_I(k) \in I \\ 0 & \text{if } l \notin I. \end{cases}$$

Now if  $m = 0$  and the image of  $\iota_I : [1] \rightarrow [n+1]$  is  $l$ , we use  $T_l$  to denote the image of  $\mathbf{c}_I : C[0] \rightarrow C[n]$ , which is called the  $t_l$  axis of  $C[n]$ . On the other hand, if  $I = \{1, \dots, \hat{l}, \dots, n+1\}$  is a subset of  $[n+1]$  with exactly one element  $l$  excluded,  $I$  is often denoted by  $I^\vee$ . We denote the image of  $\mathbf{c}_{I^\vee}$  by  $C[n]_{I^\vee}$ , which is a hypersurface of  $C[n]$ . It is clear that for  $p : C[n] \rightarrow C$ , the inverse image  $p^{-1}(\mathbf{0})$  is the union  $\cup_{l=1}^{n+1} C[n]_{I^\vee}$ .

Now we come to the standard model  $X[n]$ . Let  $\pi : X \rightarrow C$  be the family mentioned at the beginning of this section.  $X[n]$  is defined to be a desingularization of  $X \times_C C[n] = X \times_{\mathbf{A}^1} \mathbf{A}^{n+1}$  which is constructed inductively in [7] by successively blow-ups and blow-downs. There is a  $G[n]$  action on  $X \times_C C[n]$  induced from a trivial action on  $X$  and the given  $G[n]$  action on  $C[n]$ . This group action can be canonically lifted to  $X[n]$ . The following are basic properties of  $X[n]$ .

1. There are canonical  $G[n]$ -equivariant projections  $X[n] \rightarrow X$  and  $\pi_n : X[n] \rightarrow C[n]$ .

2. The central fiber  $X[n]_0 = X[n] \times_{C[n]} \mathbf{0}$  is a semistable model of  $X_0$  with  $n+2$  irreducible components.
3. Let  $\iota_I : C[m] \rightarrow C[n]$  be a standard embedding. Then the pullback  $\iota_I^* X[n]$  is canonically isomorphic to  $X[m]$ .
4. The restriction of  $X[n]$  to the  $t_I$  axis of  $C[n]$  is a smoothing of  $X[n]_0$  along the divisor  $D_I$ .

Next we study  $X[n]$  in more details. To proceed, we first recall the notion of semistable model  $X[n]_0$  of the fiber  $X_0$  of the family  $\pi : X \rightarrow C$ . Recall that  $X_0$  is a union of two smooth varieties  $Y_1$  and  $Y_2$  gluing along an isomorphism  $\psi : D_+ \cong D_-$ , where  $D_+ \subset Y_1$  and  $D_- \subset Y_2$  are smooth divisors. Let  $D = Y_1 \cap Y_2 \subset X_0$ . Then  $D \cong D_\pm$ , and we write  $X_0 = Y_1 \cup_D Y_2$ . Let  $N_i$  be the normal bundles of  $D$  in  $Y_i$ . They are line bundles and  $N_1 \cong N_2^\vee$ . Let  $\mathbf{1}_D$  be the trivial line bundle over  $D$  and let  $E = \mathbf{1}_D \oplus N_2$ . Then  $\Delta = \mathbf{P}(E)$  is a  $\mathbf{P}^1$ -bundle over  $D$ . There are two canonical sections  $s_0$  and  $s_\infty$  of  $\Delta \rightarrow D$ , which correspond to the canonical quotient bundles  $E \rightarrow N_2$  and  $E \rightarrow \mathbf{1}_D$  respectively. Because  $\Delta$  can be regarded as a projectivization of the line bundle  $N_2$ , under the canonical injection  $N_2 \rightarrow \Delta$ ,  $s_0$  is the zero section of  $N_2$  and  $s_\infty$  the infinity section. We let  $D_- = s_0(D)$  and  $D_+ = s_\infty(D)$ . Then  $D_-$  and  $D_+$  are smooth divisors in  $\Delta$ . The normal bundle of  $D_-$  in  $\Delta$  is same as  $N_2$ , and the normal bundle of  $D_+$  in  $\Delta$  is the dual  $N_2^\vee \cong N_1$ . There is a unique  $G_m$  action on  $\Delta$  up to dilations which fixes  $D_+$  and  $D_-$ . Finally, we remark that  $\Delta$  is canonically isomorphic to  $\mathbf{P}(\mathbf{1}_D \oplus N_1)$  by the isomorphism  $N_2^\vee \cong N_1$ .

Let  $\Delta_i \cong \Delta$  be the  $i^{\text{th}}$  copy of  $\Delta$  for  $i = 1, \dots, n$ . For convenience, we use  $\Delta_0$  to denote  $Y_1$  and  $\Delta_{n+1}$  to denote  $Y_2$ . A semistable model of  $X_0$  (with length  $n$ ) is a scheme with  $n+2$  smooth irreducible components

$$X[n]_0 = \Delta_0 \cup \Delta_1 \cup \dots \cup \Delta_n \cup \Delta_{n+1},$$

which is obtained by gluing  $\Delta_i$  and  $\Delta_{i+1}$  along the smooth divisors  $D_+ \subset \Delta_i$  and  $D_- \subset \Delta_{i+1}$  for  $0 \leq i \leq n$ . Let  $D_i = \Delta_{i-1} \cap \Delta_i$  for  $1 \leq i \leq n+1$ . Clearly  $D_i \cong D$ . The group  $G[n]$  acts on  $X[n]_0$  in an obvious way.

Based on the inductive construction of  $X[n]$ , we now give a description of this desingularization  $X[n] \rightarrow X \times_C C[n]$  in terms of local defining equations around a singular point  $p \times \mathbf{0} \in X \times_C C[n]$ . We begin with the original family  $\pi : X \rightarrow C$ . There is an induced closed embedding  $X \rightarrow X \times C$  whose composition with the projections to the first and second factors are identity and  $\pi$  respectively.  $X \times C$  is a constant family over  $C$  via the projection to  $C$ . Recall that  $D \subset X_0$  is the singular loci of  $X_0$ , let  $p$  be a closed point in  $D$ . We will give a local defining equation around  $p \times \mathbf{0}$  of the above closed embedding  $X \subset X \times C$ . Since  $p$  is a nonsingular point in the smooth irreducible components  $Y_i$ , we can choose local coordinate functions  $x_1, \dots, x_d$  and  $y_1, y_2$  on  $X$ , such that the complete local ring  $\hat{\mathcal{O}}_{D,p}$  is isomorphic to  $k[[x_1, \dots, x_d]]$ , and  $\hat{\mathcal{O}}_{Y_1,p} \cong k[[x_1, \dots, x_d, y_1]]$ ,  $\hat{\mathcal{O}}_{Y_2,p} \cong k[[x_1, \dots, x_d, y_2]]$ . Since  $p$  is contained in  $X_0$ , the image  $\pi(p)$  is the distinguished point  $\mathbf{0} \in C$ . Let  $s$  be a uniformizer of  $\hat{\mathcal{O}}_{C,\mathbf{0}}$ . The defining equation of  $X$  around  $p \times \mathbf{0}$  is  $y_1 y_2 = s$ . Hence the singular fiber  $X_0$  of the family  $\pi : X \rightarrow C$  is given by  $y_1 y_2 = 0$  around  $p$  in  $X$ . In general, we have a closed embedding  $X \times_C C[n] \subset X \times C[n]$ , and the local equation of  $X \times_C C[n]$  at the point  $p \times \mathbf{0}$  is  $y_1 y_2 = t_1 \cdots t_{n+1}$ . It is obvious that the scheme  $X \times_C C[n]$  is not smooth at  $p \times \mathbf{0}$ . Now we take a look at the local behavior of the desingularization at  $p \times \mathbf{0}$ . Since  $X \times C[n]$  is nonsingular, and we have already chosen local coordinate functions  $x_1, \dots, x_d, y_1, y_2, t_1, \dots, t_{n+1}$  around the

point  $p \times \mathbf{0}$ . Let  $U \subset X \times C[n]$  be a neighborhood of  $p \times \mathbf{0}$ . Consider the product  $U \times \mathbf{P}^1 \times \cdots \times \mathbf{P}^1$  with  $n$  copies of  $\mathbf{P}^1$ . Let  $u_1^i, u_2^i$  be the homogenous coordinates of the  $i^{\text{th}}$  copy of  $\mathbf{P}^1$ . We define a closed subscheme  $\hat{U}$  of  $U \times \mathbf{P}^1 \times \cdots \times \mathbf{P}^1$  by the set of  $n+1$  equations

$$u_1^1 y_1 = u_2^1 t_1, \dots, u_1^i u_2^{i-1} = u_2^i u_1^{i-1} t_i, \dots, u_1^n u_2^{n-1} = u_2^n u_1^{n-1} t_n, u_2^n y_2 = u_1^n t_{n+1}.$$

Let  $g : \hat{U} \rightarrow U$  be the projection to the first factor. Then the image of  $g$  is  $U \cap (X \times_C C[n])$ , which is a neighborhood of  $p \times \mathbf{0}$  in  $X \times_C C[n]$ . It is direct to verify that  $\hat{U}$  is smooth and it is the required desingularization  $X[n]$  of  $X \times_C C[n]$  at  $p \times \mathbf{0}$  defined in [7].

Now we describe the  $G[n]$  action on  $X[n]$ . Recall that there is a  $G[n]$  action on  $C[n]$ , we get an action on the product  $X \times_C C[n]$  and  $X \times C[n]$  from a trivial  $G[n]$  action on  $X$ . The open subset  $X \times_C C[n]^* \subset X \times_C C[n]$  is a principal  $G[n]$ -bundle over  $X \times_C C^* = X - X_0$ , the desingularization morphism is an isomorphism and the  $G[n]$  action on  $X \times_C C[n]^*$  is free. Next, without loss of generality, we consider  $p \times \mathbf{0} \in X \times_C C[n]$ . Since we choose homogenous coordinates in each copy of  $\mathbf{P}^1$ , there is a standard  $G[n]$  action on  $\mathbf{P}^1 \times \cdots \times \mathbf{P}^1$  given by

$$(u_1^1, u_2^1; u_1^2, u_2^2; \dots; u_1^n, u_2^n)^\sigma = (\sigma_1 u_1^1, u_2^1; \sigma_2 u_1^2, u_2^2; \dots; \sigma_n u_1^n, u_2^n).$$

Hence we obtain a  $G[n]$  action on  $X \times C[n] \times \mathbf{P}^1 \times \cdots \times \mathbf{P}^1$ . Notice that the defining equations of  $\hat{U}$  are  $G[n]$  invariant, it induces a  $G[n]$  action on the closed subscheme  $\hat{U}$ , which is exactly the  $G[n]$  action on  $X[n]$ .

**1.2. The stack of expanded degeneration.** We first recall the definition of effective degeneration. Let  $S$  be a  $C$ -scheme. An effective degeneration over  $S$  is a  $C$ -morphism  $\xi : S \rightarrow C[n]$  for some  $n$ . For any  $C$ -morphism  $\xi : S \rightarrow C[n]$ , there is an associated family  $\mathcal{X}$  over  $S$ , which is the pullback of the family  $\pi_n : X[n] \rightarrow C[n]$  by the morphism  $\xi$ , i.e.  $\mathcal{X} = X[n] \times_{C[n]} S$ . Clearly, we have a natural morphism  $\mathcal{X} \rightarrow X \times_C S$ . Let  $\xi_1 : S \rightarrow C[n_1]$  and  $\xi_2 : S \rightarrow C[n_2]$  be two effective degenerations over  $S$  with  $n_1 \leq n_2$ . An effective arrow  $r : \xi_1 \rightarrow \xi_2$  consists of a standard embedding  $\iota : C[n_1] \rightarrow C[n_2]$  and a morphism  $\rho : S \rightarrow G[n_2]$ , such that  $(\iota \circ \xi_1)^\rho = \xi_2$ . We let  $\mathcal{X}_1$  and  $\mathcal{X}_2$  be families over  $S$  associated with effective degenerations  $\xi_1$  and  $\xi_2$  respectively. Then an effective arrow  $r : \xi_1 \rightarrow \xi_2$  induces an  $S$ -isomorphism of the families  $\mathcal{X}_1 \cong \mathcal{X}_2$ , which is compatible to their canonical morphisms to  $X \times_C S$ .

**Definition 1.1.** Let  $S$  be a  $C$ -scheme. Let  $(\mathcal{X}, p)$  be a pair of a family  $\mathcal{X}$  flat over  $S$  and a surjective  $S$ -morphism  $p : \mathcal{X} \rightarrow X \times_C S$ . We say  $(\mathcal{X}, p)$  is an expanded degeneration over  $S$ , if there is an open covering  $\{S_\alpha\}$  of  $S$ , such that for each  $S_\alpha$  the restriction pair  $(\mathcal{X} \times_S S_\alpha, p|_{\mathcal{X} \times_S S_\alpha})$  is isomorphic to an effective degeneration over  $S_\alpha$ .

**Lemma 1.2** ([7]). Let  $\mathcal{X}_1$  and  $\mathcal{X}_2$  be two expanded degenerations over  $S$ . Suppose  $\mathcal{X}_1$  is isomorphic to  $\mathcal{X}_2$ , then for every  $p \in S$ , there is an open neighborhood  $S_0 \subset S$  of  $p$  and morphisms  $\xi_i : S_0 \rightarrow C[n_i]$ , such that  $\mathcal{X}_i \times_S S_0$  is isomorphic to the effective degeneration  $\xi_i$ , and the induced isomorphism between  $\mathcal{X}_1 \times_S S_0$  and  $\mathcal{X}_2 \times_S S_0$  is induced from a sequence of effective arrows between  $\xi_1$  and  $\xi_2$ .

Now we can define the groupoid of expanded degenerations. Let  $\mathfrak{X}$  be a category whose objects are expanded degenerations  $(\mathcal{X}, p)$ . Let  $(\mathcal{X}_1, p_1)$  and  $(\mathcal{X}_2, p_2)$  be two expanded degenerations over  $S_1$  and  $S_2$  respectively in  $\mathfrak{X}$ . An arrow  $r : (\mathcal{X}_1, p_1) \rightarrow$

$(\mathcal{X}_2, p_2)$  consists of a  $C$ -morphism  $S_1 \rightarrow S_2$  and an  $S_1$ -isomorphism  $\mathcal{X}_1 \rightarrow \mathcal{X}_2 \times_{S_2} S_1$  compatible to their canonical morphisms to  $X \times_C S_1$ . There is a functor  $F : \mathfrak{X} \rightarrow \text{Sch}/C$  sends an expanded degeneration  $(\mathcal{X}, p)$  to the base scheme  $S$  of the family  $\mathcal{X}$ . Then  $\mathfrak{X}$  together with the functor  $F$  is a groupoid over  $C$ .

**Proposition 1.3** ([7]). *The groupoid  $\mathfrak{X}$  is a stack over  $C$ .*

In the sequel, for any  $C$ -scheme  $S$ , we use  $\mathfrak{X}(S)$  to denote the groupoid of expanded degenerations over  $S$ . In particular,  $\mathfrak{X}_0$  is the closed substack of semistable models of  $X_0$ .

## 2. THE MODULI STACK OF STABLE PERFECT IDEAL SHEAVES

In this section, we first introduce the notion of perfect ideal sheaves on  $X[n]_0$ . Indeed, there are various equivalent definitions of perfectness which are convenient for different applications. We also introduce the notion of relativeness of an ideal to a smooth divisor. Next, we define the stack  $\mathfrak{I}_{X/C}^\Gamma$  of stable perfect ideal sheaves and show that it is a separated and proper Deligne-Mumford stack. Lemma 2.29 is essential in the proof of properness. To avoid distraction by the technical details, we postpone the proof of this lemma to the last section.

**2.1. Perfect ideal sheaves.** Let  $X[n]_0 = Y_1 \cup \Delta_1 \cdots \cup \Delta_n \cup Y_2$  be a semistable model of  $X_0$ . The singular locus of  $X[n]_0$  is a disjoint union  $D_+ \sqcup D_1 \sqcup \cdots \sqcup D_{n-1} \sqcup D_-$  with  $D_k = \Delta_k \cap \Delta_{k+1}$ . For convenience, we use  $\Delta_0$  to denote  $Y_1$  and  $\Delta_{n+1}$  to denote  $Y_2$ . Likewise, we let  $D_0 = D_+$  and  $D_n = D_-$ .

Let  $W$  be a Noetherian scheme and  $V \subset W$  be a closed subscheme. For any ideal sheaf  $\mathcal{I}$  on  $W$ , there is a canonical injective homomorphism  $\mathcal{I} \rightarrow \mathcal{O}_W$ . Take tensorization with  $\mathcal{O}_V$ , we obtain a homomorphism of  $\mathcal{O}_V$ -modules  $\mathcal{I} \otimes \mathcal{O}_V \rightarrow \mathcal{O}_V$  which is referred to as the canonical homomorphism.

**Definition 2.1.** *Let  $\mathcal{I}$  be an ideal sheaf on  $X[n]_0$ . We say  $\mathcal{I}$  is perfect along  $D_k$  if the canonical homomorphism  $\mathcal{I} \otimes \mathcal{O}_{D_k} \rightarrow \mathcal{O}_{D_k}$  is injective.  $\mathcal{I}$  is perfect if it is perfect along all  $D_k$  for  $k = 0, \dots, n$ .*

Let  $p$  be a closed point in  $X[n]_0$ . In the sequel, we use  $R_p$  to denote the local ring  $\mathcal{O}_{X[n]_0, p}$ . The following is an equivalent definition of perfectness.

**Lemma 2.2.** *Let  $\mathcal{I}_Z$  be an ideal sheaf of a closed subscheme  $Z \subset X[n]_0$ . Then  $\mathcal{I}_Z$  is perfect along  $D_k$  if and only if  $\text{Tor}_{R_p}^1(\mathcal{O}_{Z, p}, \mathcal{O}_{D_k, p}) = 0$  for all  $p \in D_k$ .*

Since completion is an exact functor on the category of finitely generated modules over a Noetherian ring, we get the following

**Lemma 2.3.**  *$\mathcal{I}_Z$  is perfect along  $D_k$  if and only if the canonical homomorphism  $\hat{\mathcal{I}}_p \otimes \hat{\mathcal{O}}_{D_k, p} \rightarrow \hat{\mathcal{O}}_{D_k, p}$  is injective for all closed points  $p \in D_k$ , or equivalently,  $\text{Tor}_{\hat{R}_p}^1(\hat{\mathcal{O}}_{Z, p}, \hat{\mathcal{O}}_{D_k, p}) = 0$  for all  $p \in D_k$ .*

**Lemma 2.4.** *Let  $\mathcal{I}$  be an ideal sheaf of a closed subscheme  $Z \subset X_0 = Y_1 \cup_D Y_2$ . Then  $\text{Tor}_{R_p}^1(\mathcal{O}_{Z, p}, \mathcal{O}_{D, p}) = 0$  if and only if  $\text{Tor}_{R_p}^1(\mathcal{O}_{Z, p}, \mathcal{O}_{Y_i, p}) = 0$  for both  $i = 1, 2$ .*

*Proof.* First we show the “only if” part. Since there is an exact sequence

$$0 \longrightarrow \mathcal{O}_{X_0}(-D) \longrightarrow \mathcal{O}_{X_0} \longrightarrow \mathcal{O}_D \longrightarrow 0,$$

and an isomorphism  $\mathcal{O}_{X_0}(-D) \cong \mathcal{O}_{X_0}(-Y_1) \oplus \mathcal{O}_{X_0}(-Y_2)$ , we obtain an exact sequence

$$0 \longrightarrow \mathcal{O}_{X_0}(-Y_1) \oplus \mathcal{O}_{X_0}(-Y_2) \xrightarrow{f} \mathcal{O}_{X_0} \longrightarrow \mathcal{O}_D \longrightarrow 0.$$

Consider the sequence of stalks at  $p$  and take tensorization with  $\mathcal{O}_{Z,p}$ , we obtain

$$\mathrm{Tor}_{R_p}^1(\mathcal{O}_{Z,p}, \mathcal{O}_{D,p}) \longrightarrow \mathcal{O}_{X_0}(-Y_1)_p \otimes \mathcal{O}_{Z,p} \oplus \mathcal{O}_{X_0}(-Y_2)_p \otimes \mathcal{O}_{Z,p} \xrightarrow{f \otimes 1} \mathcal{O}_{Z,p} \longrightarrow \cdots.$$

Since  $\mathrm{Tor}_{R_p}^1(\mathcal{O}_{Z,p}, \mathcal{O}_{D,p}) = 0$ ,  $f \otimes 1$  is injective. Notice that there are exact sequences

$$0 \longrightarrow \mathcal{O}_{X_0}(-Y_i) \xrightarrow{f_i} \mathcal{O}_{X_0} \longrightarrow \mathcal{O}_{Y_i} \longrightarrow 0,$$

and  $f = f_1 + f_2$ , the injectivity of  $f \otimes 1$  implies that both  $f_i \otimes 1$  are also injective, which is equivalent to  $\mathrm{Tor}_{R_p}^1(\mathcal{O}_{Z,p}, \mathcal{O}_{Y_i,p}) = 0$  for both  $i = 1, 2$ .

Next, we prove the “if” part. Suppose that  $\mathrm{Tor}_{R_p}^1(\mathcal{O}_{Z,p}, \mathcal{O}_{Y_i,p}) = 0$  for  $i = 1, 2$ , then  $\mathrm{Tor}_{R_p}^1(\hat{\mathcal{O}}_{Z,p}, \hat{\mathcal{O}}_{Y_i,p}) = 0$ . We will show that the induced homomorphism

$$\hat{f} \otimes 1 : \hat{\mathcal{O}}_{X_0}(-Y_1)_p \otimes \hat{\mathcal{O}}_{Z,p} \oplus \hat{\mathcal{O}}_{X_0}(-Y_2)_p \otimes \hat{\mathcal{O}}_{Z,p} \xrightarrow{\hat{f} \otimes 1} \hat{\mathcal{O}}_{Z,p}$$

is injective, which is equivalent to  $\mathrm{Tor}_{R_p}^1(\hat{\mathcal{O}}_{Z,p}, \hat{\mathcal{O}}_{D_k,p}) = 0$  and thus the result follows. Let  $a \in \hat{\mathcal{O}}_{X_0}(-Y_1)_p$  and  $b \in \hat{\mathcal{O}}_{X_0}(-Y_2)_p$  such that the image of  $a \otimes 1 + b \otimes 1$  via  $\hat{f} \otimes 1$  is zero in  $\hat{\mathcal{O}}_{Z,p}$ . Since  $\hat{R}_p \cong \hat{\mathcal{O}}_{D,p}[[y_1, y_2]]/(y_1 y_2)$  and  $\hat{\mathcal{O}}_{X_0}(-Y_1)_p = (y_2)$  and  $\hat{\mathcal{O}}_{X_0}(-Y_2)_p = (y_1)$  as ideals in  $\hat{R}_p$ , we have  $a = y_2 a'$  and  $b = y_1 b'$  for some  $a', b' \in \hat{R}_p$ . Now  $\hat{f} \otimes 1$  sends  $a \otimes 1 + b \otimes 1$  to  $y_2 a' + y_1 b'$  that is zero in  $\hat{\mathcal{O}}_{Z,p}$ . It implies that  $y_2 a' + y_1 b' \in \hat{\mathcal{I}}_p$ . Multiplied by  $y_2$ , and since  $y_1 y_2 = 0$ , we get  $y_2^2 a' \in \hat{\mathcal{I}}_p$ . Because  $\mathrm{Tor}_{R_p}^1(\hat{\mathcal{O}}_{Z,p}, \hat{\mathcal{O}}_{Y_1,p}) = 0$ , the canonical homomorphism  $\hat{\mathcal{I}}_p \otimes \hat{\mathcal{O}}_{Y_1,p} \rightarrow \hat{\mathcal{O}}_{Y_1,p}$  is injective, and it sends  $y_2^2 a' \otimes 1$  to  $y_2^2 a'$  that is zero in  $\hat{\mathcal{O}}_{Y_1,p}$ . It implies that  $a' \in \hat{\mathcal{I}}_p$ . Thus  $a \otimes 1 = y_2 a' \otimes 1 = y_2 \otimes a'$  is zero in  $\hat{\mathcal{O}}_{X_0}(-Y_1)_p \otimes \hat{\mathcal{O}}_{Z,p}$ . Likewise,  $b \otimes 1 = 0$ . Therefore  $a \otimes 1 + b \otimes 1 = 0$  and the injectivity of  $\hat{f} \otimes 1$  follows.  $\square$

**Remark 2.5.** The “only if” part of this lemma is true for any coherent sheaf  $\mathcal{F}$  on  $X_0$ . Using the exact sequences in the proof, it is straightforward that for  $k > 1$ ,  $\mathrm{Tor}_{R_p}^k(\mathcal{F}, \mathcal{O}_{D,p}) = 0$  if and only if  $\mathrm{Tor}_{R_p}^k(\mathcal{F}, \mathcal{O}_{Y_i,p}) = 0$  for both  $i = 1, 2$ .

**Corollary 2.6.** Let  $\mathcal{I}$  be an ideal sheaf on  $X[n]_0$ . Then  $\mathcal{I}$  is perfect if and only if for every irreducible component  $\Delta_k$  of  $X[n]_0$ , the canonical homomorphism  $\mathcal{I} \otimes \mathcal{O}_{\Delta_k} \rightarrow \mathcal{O}_{\Delta_k}$  is injective.

Next we use another point of view to study perfectness. In what follows, a free module always means a finitely generated free module. We begin by stating a fact in commutative algebra.

**Lemma 2.7.** Let  $A$  be a regular local ring. Let  $M$  be a finitely generated torsion free  $A$ -module. Suppose  $x \in A$  is not a unit and  $A' = A/(x)$ . If  $M \otimes_A A'$  is a free  $A'$ -module, then  $M$  is a free  $A$ -module.

**Lemma 2.8.** Let  $Z \subset X_0$  be a closed subscheme and  $\dim X_0 = d$ . Let  $p$  be a closed point in  $D$  and recall that  $R_p = \mathcal{O}_{X_0,p}$ . Then the  $R_p$ -algebra  $\mathcal{O}_{Z,p}$  admits a length  $d - 1$  resolution of free  $R_p$ -modules if and only if  $\mathrm{Tor}_{R_p}^i(\mathcal{O}_{Z,p}, \mathcal{O}_{D,p}) = 0$  for all  $i > 0$ .



*Proof.* First we prove the “if” part. In fact, this part is true for any coherent sheaf  $\mathcal{F}$  on  $X_0$ . Now we prove this general result. Since  $\mathcal{F}_p$  is finitely generated, there exists an exact sequence

$$0 \longrightarrow K \xrightarrow{g} E_{d-1} \longrightarrow \cdots \longrightarrow E_0 \longrightarrow \mathcal{F}_p \longrightarrow 0,$$

such that all  $E_i$  are free  $R_p$ -modules. Now we show that  $K$  and  $\text{coker } g$  are also free  $R_p$ -modules, which implies that  $\mathcal{F}_p$  admits a length  $d-1$  free resolution. Since  $\text{Tor}_{R_p}^i(\mathcal{F}_p, \mathcal{O}_{D,p}) = 0$  for all  $i > 0$ , followed by the argument of “only if” part in Lemma 2.4, we get  $\text{Tor}_{R_p}^i(\mathcal{F}_p, \mathcal{O}_{Y_1,p}) = 0$  and  $\text{Tor}_{R_p}^i(\mathcal{F}_p, \mathcal{O}_{Y_2,p}) = 0$  for  $i > 0$ . It follows that by taking tensorization with  $\mathcal{O}_{Y_1,p}$ ,  $\mathcal{O}_{Y_2,p}$  and  $\mathcal{O}_{D,p}$  on the above exact sequence respectively, we obtain exact sequences

$$0 \longrightarrow K^+ \xrightarrow{g_1} E_{d-1}^+ \longrightarrow \cdots \longrightarrow E_0^+ \longrightarrow \mathcal{F}_p^+ \longrightarrow 0.$$

$$0 \longrightarrow K^- \xrightarrow{g_2} E_{d-1}^- \longrightarrow \cdots \longrightarrow E_0^- \longrightarrow \mathcal{F}_p^- \longrightarrow 0.$$

$$0 \longrightarrow K^D \xrightarrow{g'} E_{d-1}^D \longrightarrow \cdots \longrightarrow E_0^D \longrightarrow \mathcal{F}_p^D \longrightarrow 0.$$

where  $E_j^+ = E_j \otimes \mathcal{O}_{Y_1,p}$ ,  $E_j^- = E_j \otimes \mathcal{O}_{Y_2,p}$  and  $E_j^D = E_j \otimes \mathcal{O}_{D,p}$ . Therefore we have the following commutative diagram

$$\begin{array}{ccccccc} & & 0 & & 0 & & \cdots & & 0 \\ & & \downarrow & & \downarrow & & & & \downarrow \\ 0 & \longrightarrow & K & \longrightarrow & E_{d-1} & \longrightarrow & \cdots & \longrightarrow & \mathcal{F}_p & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & & & \downarrow & & \\ 0 & \longrightarrow & K^+ \oplus K^- & \longrightarrow & E_{d-1}^+ \oplus E_{d-1}^- & \longrightarrow & \cdots & \longrightarrow & \mathcal{F}_p^+ \oplus \mathcal{F}_p^- & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & & & \downarrow & & \\ 0 & \longrightarrow & K^D & \longrightarrow & E_{d-1}^D & \longrightarrow & \cdots & \longrightarrow & \mathcal{F}_p^D & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & & & \downarrow & & \\ & & 0 & & 0 & & \cdots & & 0 \end{array}$$

Since  $\mathcal{O}_{Y_1,p}$  and  $\mathcal{O}_{Y_2,p}$  are regular rings with dimension  $d$ ,  $\mathcal{F}_p^+$  and  $\mathcal{F}_p^-$  have length  $d$  free resolutions. It follows that  $K^+$  and  $K^-$  are free modules, which implies that  $K$  is also a free module. Let  $Q$  be the cokernel of  $g$ . We will show that  $Q$  is a free module. Since  $\mathcal{O}_{D,p}$  is regular with dimension  $d-1$ ,  $Q \otimes \mathcal{O}_{D,p} = \text{coker } g'$  is a free module. Notice that for  $i = 1, 2$ ,  $Q \otimes \mathcal{O}_{Y_i,p} = \text{coker } g_i$  are torsion free, by Lemma 2.7,  $Q \otimes \mathcal{O}_{Y_i,p}$  are both free modules. Hence  $Q$  is also a free module.

Next, we prove the “only if” part. We also start with a general coherent sheaf  $\mathcal{F}$  on  $X_0$ . Suppose there is a length  $d-1$  free resolution of  $\mathcal{F}_p$

$$0 \longrightarrow E_{d-1} \longrightarrow \cdots \longrightarrow E_0 \longrightarrow \mathcal{F}_p \longrightarrow 0,$$

then  $\text{Tor}_{R_p}^i(\mathcal{F}_p, M) = 0$  for all  $i > d$  and all  $R_p$ -modules  $M$ . In particular, we let  $M = \mathcal{O}_{Y_1,p}$ . Since there is an exact sequence

$$\longrightarrow R_p \longrightarrow \cdots \longrightarrow R_p \xrightarrow{f_3} R_p \xrightarrow{f_2} R_p \xrightarrow{f_1} R_p \longrightarrow \mathcal{O}_{Y_1,p} \longrightarrow 0,$$

with  $f_1 = f_3 = \cdots = f_{2k-1}$  and  $f_2 = f_4 = \cdots = f_{2k}$  for any integer  $k > 0$ . It is clear that  $\text{im } f_1 = \mathcal{O}_{Y_2,p}$  and  $\text{im } f_2 = \mathcal{O}_{Y_1,p}$ . Since

$$\text{Tor}_{R_p}^i(\mathcal{F}_p, \mathcal{O}_{Y_1,p}) = \text{Tor}_{R_p}^{i-1}(\mathcal{F}_p, \text{im } f_1) = \cdots = \text{Tor}_{R_p}^1(\mathcal{F}_p, \text{im } f_{i-1})$$

for any  $i > 1$ , and recall that  $\text{Tor}_{R_p}^i(\mathcal{F}_p, \mathcal{O}_{Y_1,p}) = 0$  for all  $i > d$ , we obtain  $\text{Tor}_{R_p}^i(\mathcal{F}_p, \mathcal{O}_{Y_1,p}) = 0$  and  $\text{Tor}_{R_p}^i(\mathcal{F}_p, \mathcal{O}_{Y_2,p}) = 0$  for all  $i > 0$ . Let  $\mathcal{F} = \mathcal{O}_Z$  for a closed subscheme  $Z \subset X_0$ . Then by Lemma 2.4,  $\text{Tor}_{R_p}^i(\mathcal{O}_{Z,p}, \mathcal{O}_{D,p}) = 0$  for all  $i > 0$ .  $\square$

The argument in this proof also shows that  $\mathcal{O}_{Z,p}$  admits a length  $d - 1$  free resolution if and only if it admits a finite length free resolution.

**Corollary 2.9.** *Let  $\mathcal{I}$  be an ideal sheaf on  $X[n]_0$ . Then  $\mathcal{I}$  is perfect if and only if for every closed point  $p \in X[n]_0$ , the ideal  $\mathcal{I}_p$  admits a finite length resolution of free modules.*

In fact, this corollary can be regarded as another definition of perfectness.

**2.2. Relativeness of ideal sheaves.** The notion of relativeness of ideal sheaves is closely related to perfectness.

**Definition 2.10.** *Let  $Y$  be a smooth variety and  $D \subset Y$  a smooth divisor. An ideal sheaf  $\mathcal{I} \subset \mathcal{O}_Y$  is called relative with respect to  $D$  if the canonical homomorphism  $\mathcal{I} \otimes \mathcal{O}_D \rightarrow \mathcal{O}_D$  is injective.*

Relativeness is a local property. We say  $\mathcal{I}$  is relative at a closed point  $p \in D$  if the canonical homomorphism of stalks  $\mathcal{I}_p \otimes \mathcal{O}_{D,p} \rightarrow \mathcal{O}_{D,p}$  is injective. Then  $\mathcal{I}$  is relative with respect to  $D$  if and only if it is relative at every closed point  $p \in D$ .

Similar to perfectness, the following two lemmas are equivalent definition of relativeness.

**Lemma 2.11.** *Let  $\mathcal{I}$  be an ideal sheaf on  $Y$ . Then  $\mathcal{I}$  is relative at  $p \in D$  if and only if the canonical homomorphism  $\hat{\mathcal{I}}_p \otimes \hat{\mathcal{O}}_{D,p} \rightarrow \hat{\mathcal{O}}_{D,p}$  is injective.*

**Lemma 2.12.** *Let  $\mathcal{I}_Z$  be an ideal sheaf of a closed subscheme  $Z \subset Y$ . Then  $\mathcal{I}_Z$  is relative at a point  $p \in D$  if and only if  $\text{Tor}_{\mathcal{O}_{Y,p}}^1(\mathcal{O}_{Z,p}, \mathcal{O}_{D,p}) = 0$ , or equivalently,  $\text{Tor}_{\hat{\mathcal{O}}_{Y,p}}^1(\hat{\mathcal{O}}_{Z,p}, \hat{\mathcal{O}}_{D,p}) = 0$ .*

The relation between relativeness and perfectness is straightforward. Let  $\mathcal{I}$  be an ideal sheaf on  $X[n]_0$ . If  $\mathcal{I}$  is perfect, then by Corollary 2.6,  $\mathcal{I} \otimes \mathcal{O}_{\Delta_k}$  is an ideal sheaf on  $\Delta_k$ , and it is clear that  $\mathcal{I} \otimes \mathcal{O}_{\Delta_k}$  is relative with respect to the distinguished divisors  $D_-$  and  $D_+$  of  $\Delta_k$  for all  $k$ . Note that for  $\Delta_0$  and  $\Delta_{n+1}$ , there is only one distinguished divisor. Conversely, we have

**Lemma 2.13.** *For every  $k$ , let  $\mathcal{I}_k$  be an ideal sheaf on  $\Delta_k$ . Suppose  $\mathcal{I}_k$  is relative with respect to the distinguished divisors  $D_-$  and  $D_+$  of  $\Delta_k$  and  $\mathcal{I}_k \otimes \mathcal{O}_{D_k} \cong \mathcal{I}_{k+1} \otimes \mathcal{O}_{D_k}$ . Then there is a perfect ideal sheaf  $\mathcal{I}$  on  $X[n]_0$  such that  $\mathcal{I} \otimes \mathcal{O}_{\Delta_k} \cong \mathcal{I}_k$ .*

*Proof.* It suffices to consider the case of ideal sheaves on  $X_0$ . Given ideal sheaves  $\mathcal{I}_1$  and  $\mathcal{I}_2$  on  $Y_1$  and  $Y_2$  respectively, since  $\mathcal{I}_1 \otimes \mathcal{O}_D \cong \mathcal{I}_2 \otimes \mathcal{O}_D$ , we let  $\mathcal{I}^D = \mathcal{I}_1 \otimes \mathcal{O}_D$  and  $\phi_i : \mathcal{I}_i \rightarrow \mathcal{I}^D$  be the restriction homomorphisms. Let  $\phi : \mathcal{I}_1 \oplus \mathcal{I}_2 \rightarrow \mathcal{I}^D$  be the surjective homomorphism defined by  $\phi(a, b) = \phi_1(a) - \phi_2(b)$  for  $a \in \mathcal{I}_1$  and  $b \in \mathcal{I}_2$ .

We will show that  $\ker \phi$  is a perfect ideal sheaf on  $X_0$  whose restrictions on  $Y_i$  are exactly  $\mathcal{I}_i$ .

From the commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \ker \phi & \longrightarrow & \mathcal{I}_1 \oplus \mathcal{I}_2 & \longrightarrow & \mathcal{I}^D \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \mathcal{O}_{X_0} & \longrightarrow & \mathcal{O}_{Y_1} \oplus \mathcal{O}_{Y_2} & \longrightarrow & \mathcal{O}_D \longrightarrow 0 \end{array}$$

we see that  $\ker \phi$  is an ideal sheaf on  $X_0$ . Let  $\mathcal{I} = \ker \phi$ . If  $\mathcal{I}|_{Y_i}$  has a subsheaf  $\mathcal{T}_i$  supported on a subscheme of codimension at least one, we let  $(\mathcal{I}|_{Y_i})^p = \text{coker}(\mathcal{T}_i \rightarrow \mathcal{I}|_{Y_i})$ . Since  $\phi_i$  are surjective,  $(\mathcal{I}|_{Y_i})^p \cong \mathcal{I}_i$  and the homomorphism  $\mathcal{I} \rightarrow \mathcal{I}_1 \oplus \mathcal{I}_2$  is equivalent to  $\mathcal{I} \rightarrow (\mathcal{I}|_{Y_1})^p \oplus (\mathcal{I}|_{Y_2})^p$ . Since there is an exact sequence

$$0 \longrightarrow \mathcal{I} \longrightarrow \mathcal{I}|_{Y_1} \oplus \mathcal{I}|_{Y_2} \longrightarrow \mathcal{I}|_D \longrightarrow 0,$$

and  $\mathcal{I}$  is torsion free, the intersection of images of  $\mathcal{T}_i$  in  $\mathcal{I}|_D$  is zero. Hence if  $\mathcal{T}_i$  are not both zero, we get a contradiction from  $\mathcal{I}^D \cong \text{coker}(\mathcal{I} \rightarrow \mathcal{I}|_D)$ . Since  $\mathcal{I}|_{Y_i}$  are both torsion free, by Corollary 2.6,  $\mathcal{I}$  is perfect.  $\square$

Next we introduce a criterion for relativeness of an ideal sheaf. To begin with, we introduce some notation. As usual, let  $Y$  be a smooth variety with a smooth divisor  $D$ . Let  $p$  be a closed point in  $D$ . Since there is a surjective homomorphism  $\pi : \hat{\mathcal{O}}_{Y,p} \rightarrow \hat{\mathcal{O}}_{D,p}$  between complete local rings, we can choose local coordinates  $x_1, \dots, x_d, y$  in  $\hat{\mathcal{O}}_{Y,p}$ , such that  $\pi(y) = 0$  in  $\hat{\mathcal{O}}_{D,p}$ , and  $\hat{\mathcal{O}}_{Y,p} = k[[y, x_1, \dots, x_d]]$ ,  $\hat{\mathcal{O}}_{D,p} = k[[\bar{x}_1, \dots, \bar{x}_d]]$ , where  $\bar{x}_i = \pi(x_i)$ .

**Proposition 2.14.** *Let  $\mathcal{I}$  be an ideal sheaf on  $Y$ . Then the following statements are equivalent:*

- (1)  $\mathcal{I}$  is relative at  $p \in D$ .
- (2) Let  $u \in \hat{\mathcal{O}}_{Y,p}$ . If  $yu \in \hat{\mathcal{I}}_p$ , then  $u \in \hat{\mathcal{I}}_p$ .

*Proof.* Let  $Z \subset X$  be the closed subscheme defined by  $\mathcal{I}$ . Then there is a canonical exact sequence

$$0 \longrightarrow \mathcal{I} \longrightarrow \mathcal{O}_X \longrightarrow \mathcal{O}_Z \longrightarrow 0.$$

Take tensorization with  $\mathcal{O}_D$ , we obtain an exact sequence

$$\mathcal{I} \otimes \mathcal{O}_D \longrightarrow \mathcal{O}_D \longrightarrow \mathcal{O}_Z \otimes \mathcal{O}_D \longrightarrow 0.$$

The kernel of the homomorphism  $\mathcal{O}_D \rightarrow \mathcal{O}_Z \otimes \mathcal{O}_D$  is an ideal sheaf of  $\mathcal{O}_D$ , which defines the closed subscheme  $Z \cap D \subset D$ . This ideal sheaf is exactly the image of  $\mathcal{I}$  under the surjective homomorphism  $\mathcal{O}_X \rightarrow \mathcal{O}_D$ , which is denoted by  $\mathcal{I}\mathcal{O}_D$ . The relativeness of  $\mathcal{I}$  at  $p \in D$  is equivalent to the following condition: the canonical surjective homomorphism  $\hat{\mathcal{I}}_p \otimes \hat{\mathcal{O}}_{D,p} \rightarrow \hat{\mathcal{I}}_p \mathcal{O}_{D,p}$  is an isomorphism, or equivalently,  $\hat{\mathcal{I}}_p \otimes \hat{\mathcal{O}}_{D,p} \rightarrow \hat{\mathcal{I}}_p \hat{\mathcal{O}}_{D,p}$  is an isomorphism.

We first show that (2) implies (1). Since  $\pi : \hat{\mathcal{O}}_{Y,p} \rightarrow \hat{\mathcal{O}}_{D,p}$  is surjective, every element in  $\hat{\mathcal{I}}_p \otimes \hat{\mathcal{O}}_{D,p}$  can be written as  $w \otimes 1$  for some  $w \in \hat{\mathcal{I}}_p$ . The image of  $w \otimes 1$  in  $\hat{\mathcal{I}}_p \hat{\mathcal{O}}_{D,p}$  is  $\pi(w)$ . Now suppose  $\pi(w) = 0$ , we need to show that  $w \otimes 1 = 0$ , which would imply that  $\hat{\mathcal{I}}_p \otimes \hat{\mathcal{O}}_{D,p} \rightarrow \hat{\mathcal{I}}_p \hat{\mathcal{O}}_{D,p}$  is an isomorphism. Because  $D$  is defined by  $y = 0$  locally around  $p$ ,  $\pi(w) = 0$  implies that  $w = yu$  for some  $u \in \hat{\mathcal{O}}_{Y,p}$ . Now

$yu = w \in \hat{\mathcal{I}}_p$  and by (2), we have  $u \in \hat{\mathcal{I}}_p$ . Therefore  $w \otimes 1 = yu \otimes 1 = u \otimes y = 0$  in  $\hat{\mathcal{I}}_p \otimes \hat{\mathcal{O}}_{D,p}$ .

Next we show that (1) implies (2). By lemma 2.12, the relativeness of  $\mathcal{I}$  at  $p$  is equivalent to  $\text{Tor}_{\hat{\mathcal{O}}_{X,p}}^1(\hat{\mathcal{O}}_{Z,p}, \hat{\mathcal{O}}_{D,p}) = 0$ . It implies that if we tensor  $\hat{\mathcal{O}}_{Z,p}$  on the exact sequence

$$0 \longrightarrow (y) \longrightarrow \hat{\mathcal{O}}_{X,p} \longrightarrow \hat{\mathcal{O}}_{D,p} \longrightarrow 0,$$

then we get an exact sequence

$$0 \longrightarrow (y) \otimes \hat{\mathcal{O}}_{Z,p} \longrightarrow \hat{\mathcal{O}}_{Z,p} \longrightarrow \hat{\mathcal{O}}_{D,p} \otimes \hat{\mathcal{O}}_{Z,p} \longrightarrow 0.$$

Let  $yu$  be an element in  $\hat{\mathcal{I}}_p$  and  $\overline{yu}$  be its image in  $\hat{\mathcal{O}}_{Z,p}$  under the homomorphism  $\hat{\mathcal{O}}_{X,p} \rightarrow \hat{\mathcal{O}}_{Z,p}$ . Then  $\overline{yu} = 0$ . Notice that  $\overline{yu}$  is the image of  $y \otimes \bar{u}$  via the injective homomorphism  $(y) \otimes \hat{\mathcal{O}}_{Z,p} \rightarrow \hat{\mathcal{O}}_{Z,p}$ , it implies that  $y \otimes \bar{u} = 0$  in  $(y) \otimes \hat{\mathcal{O}}_{Z,p}$ . It follows that  $\bar{u} = 0$  in  $\hat{\mathcal{O}}_{Z,p}$ . Therefore  $u$  is contained in  $\hat{\mathcal{I}}_p$ .  $\square$

As a corollary, we obtain the following complete description of relative ideal sheaves  $\mathcal{I}$  provided that the subscheme  $Z$  defined by  $\mathcal{I}$  is pure, that is, the support of any subsheaf of  $\mathcal{O}_Z$  has the same dimension as  $Z$ .

**Corollary 2.15.** *Let  $Y$  be a smooth variety with a smooth divisor  $D \subset Y$ . Let  $\mathcal{I}_Z \subset \mathcal{O}_Y$  be an ideal sheaf of a closed subscheme  $Z$  of pure dimension. Then  $\mathcal{I}_Z$  is relative to  $D$  if and only if the intersection of  $Z$  and  $D$  is proper, i.e.  $\dim(Z \cap D) = \dim Z - 1$ .*

In particular, when the dimension of  $Z$  is 0 or 1, we obtain

**Corollary 2.16.** *Let  $Z \subset Y$  be a zero dimensional closed subscheme. Then the ideal sheaf  $\mathcal{I}_Z \subset \mathcal{O}_Y$  is relative to  $D$  if and only if  $Z \cap D = \emptyset$ .*

**Corollary 2.17.** *Let  $Z \subset Y$  be a pure closed subscheme of dimension one. The ideal sheaf  $\mathcal{I}_Z \subset \mathcal{O}_Y$  is relative to  $D$  if and only if every irreducible component of  $Z$  is not contained in  $D$ .*

**2.3. Families of perfect ideal sheaves.** We now turn to families of perfect ideal sheaves. To proceed, we need the following fact. Let  $\mathcal{X}/S$  be an expanded degeneration. Then for every closed point  $s \in S$ , the fibre  $\mathcal{X}_s$  is a reduced scheme. In fact,  $\mathcal{X}_s$  is either a smooth fibre of  $X/C$  or a semistable model  $X[n]_0$  for some  $n$ .

We defined perfectness for ideal sheaves on semistable models of  $X_0$ . Let  $\mathcal{X}_s$  be a smooth fibre in an expanded degeneration  $\mathcal{X}/S$ . We make the convention that every ideal sheaf on  $\mathcal{X}_s$  is perfect. This is suitable because the singular locus of  $\mathcal{X}_s$  is empty. If we use Corollary 2.9 as the definition of perfectness, then this convention is an immediate fact.

Let  $\iota_t : X_t \rightarrow X$  be the closed embedding of the fibre  $X_t$  of the family  $X/C$ . We fix a homology class  $\Gamma \in H_*(X, \mathbf{Z})$ . For any  $t \in C$ , one can always find a class  $\Gamma_t \in H_*(X_t, \mathbf{Z})$  such that  $\iota_t(\Gamma_t) = \Gamma$ . We will define families of perfect ideal sheaves  $\mathcal{I}$  on  $\mathcal{X}$  such that on the fibre  $\mathcal{X}_t$ , the Poincare dual of the Chern class  $c(\mathcal{I}_t)$  is  $\Gamma_t$ . If  $\mathcal{X}_t$  is smooth, everything is well defined. When  $\mathcal{X}_t$  is singular, we proceed as follows. First,  $\mathcal{X}_t$  is isomorphic to  $X[n]_0$  for some  $n$ . By Lemma 2.9, one can define Chern class  $c(\mathcal{I}) \in H^*(X[n]_0, \mathbf{Z})$  for a perfect ideal sheaves  $\mathcal{I}$ . Let  $\mu_i : \Delta_i \rightarrow X[n]_0$  be the closed embedding of the  $i^{\text{th}}$  irreducible component. It induces a homomorphism  $\mu_i^* : H^*(X[n]_0, \mathbf{Z}) \rightarrow H^*(\Delta_i, \mathbf{Z})$ . Let  $\omega_i = \mu_i^*(c)$  and  $\gamma_i$  be the Poincare dual of  $\omega_i$ .

Let  $\gamma = \sum \mu_*(\gamma_i)$  be a homology class in  $H_*(X[n]_0, \mathbf{Z})$ . Sometimes we use  $\gamma(\mathcal{I})$  to stress the dependence of  $\gamma$  to  $\mathcal{I}$ . Let  $\Gamma_0$  be the mentioned fixed homology class on  $X_0$ . Now we say  $\gamma \sim \Gamma_0$  if  $p_*(\gamma) = \Gamma_0$  where  $p : X[n]_0 \rightarrow X_0$  is the canonical projection.

**Definition 2.18.** *A perfect ideal sheaf  $\mathcal{I}$  on  $\mathcal{X}_s$  has type  $\Gamma$  for smooth  $\mathcal{X}_s \cong X_t$  if the Poincare dual of the Chern class  $c(\mathcal{I})$  is  $\Gamma_t$ , and for singular  $\mathcal{X}_t = X[n]_0$ , if  $\gamma(\mathcal{I}) \sim \Gamma_0$ .*

**Definition 2.19.** *A family of perfect ideal sheaves of type  $\Gamma$  on an expanded degeneration  $\mathcal{X}/S$  is an ideal sheaf  $\mathcal{I}$  of a closed subscheme  $Z \subset \mathcal{X}$ , such that  $\mathcal{O}_Z$  is flat over  $S$ , and for every closed point  $s \in S$ ,  $\mathcal{I}_s = \mathcal{I} \otimes_{\mathcal{O}_S} k_s$  is a perfect ideal sheaf on  $\mathcal{X}_s$  of type  $\Gamma$ .*

A remark is in order. If  $\mathcal{O}_Z$  is flat over  $S$ , then  $\mathcal{I}_s$  is always an ideal sheaf on  $\mathcal{X}_s$ . In the sequel, when we say a family of ideal sheaves  $\mathcal{I}$  on  $\mathcal{X}/S$ , it always means the closed subscheme  $Z$  defined by  $\mathcal{I}$  is flat over  $S$ .

**Proposition 2.20.** *Let  $\mathcal{I}$  be a family of ideal sheaves on  $\mathcal{X}/S$ . Then the locus of points  $s \in S$  such that  $\mathcal{I}_s$  is perfect is a Zariski open subset of  $S$ .*

*Proof.* Let  $U \subset S$  be the subset such that for any  $s \in U$ ,  $\mathcal{I}_s$  is perfect. Suppose  $U$  is nonempty, for otherwise it is already an open subset. Let  $s_0 \in U$  be a closed point. Then  $\mathcal{I}_{s_0}$  is perfect. We will show that there exists a Zariski open subset  $V$  containing  $s_0$  and  $V \subset U$ .

Let

$$\mathcal{E}_{d-3} \xrightarrow{f} \cdots \longrightarrow \mathcal{E}_1 \longrightarrow \mathcal{E}_0 \longrightarrow \mathcal{I} \longrightarrow 0$$

be an exact sequence of  $\mathcal{O}_{\mathcal{X}}$ -modules with  $\mathcal{E}_i$  locally free. Let  $\mathcal{K}$  be the kernel of  $f$ . Since  $\mathcal{I}$  is flat over  $S$ ,  $\mathcal{K}$  is also flat over  $S$ , and the sequence

$$0 \longrightarrow \mathcal{K} \otimes k_{s_0} \longrightarrow \mathcal{E}_{d-3} \otimes k_{s_0} \xrightarrow{f} \cdots \longrightarrow \mathcal{E}_0 \otimes k_{s_0} \longrightarrow \mathcal{I}_{s_0} \longrightarrow 0$$

is also exact. Since  $\mathcal{I}_s$  is perfect, by Corollary 2.9,  $\mathcal{K} \otimes k_{s_0}$  is a locally free sheaf on  $\mathcal{X}_{s_0}$ . The flatness of  $\mathcal{K}$  implies that  $\mathcal{K}_p$  is a free  $\mathcal{O}_{\mathcal{X},p}$ -module for every point  $p \in \mathcal{X}_{s_0}$ . Therefore there is a Zariski open subset  $V \ni s_0$  such that for any  $s \in V$ ,  $\mathcal{K} \otimes k_s$  is locally free. Apply Corollary 2.9 again, we know  $\mathcal{I}_s$  is a perfect ideal sheaf on  $\mathcal{X}_s$ . It implies that  $V \subset U$  and hence  $U$  is an open subset.  $\square$

**2.4. The stability condition.** Let  $\mathcal{I}$  be an ideal sheaf on  $X[n]_0$ . A categorical automorphism of  $\mathcal{I}$  consists of an isomorphism  $u : X[n]_0 \rightarrow X[n]_0$  and an isomorphism  $\mathcal{I} \cong u^*\mathcal{I}$  as ideal sheaves on  $X[n]_0$ . Since the automorphism group of  $X[n]_0$  is  $G[n]$ , the group of categorical automorphisms of  $\mathcal{I}$  is a subgroup of  $G[n]$ .

**Definition 2.21.** *Let  $\mathcal{I}$  be an ideal sheaf on  $X[n]_0$ .  $\mathcal{I}$  is called quasi-stable if its group of categorical automorphisms is finite.  $\mathcal{I}$  is called stable if it is perfect and quasi-stable.*

To avoid confusion, we use the word "ordinary stable" to refer to Gieseker stability of a coherent sheaf.

**Definition 2.22.** *Let  $\mathcal{I}$  be an ideal sheaf on  $X[n]_0$ . Let  $\mathcal{I}_k = \mathcal{I} \otimes \mathcal{O}_{\Delta_k}$ . Then  $\Delta_k$  is called a trivial component of  $\mathcal{I}$  if  $\mathcal{I}_k \cong \lambda^*\mathcal{I}_k$  for every dilation  $\lambda$  of  $\Delta_k$ .*

**Lemma 2.23.** *Let  $\mathcal{I}$  be an ideal sheaf on  $X[n]_0$ . Then  $\mathcal{I}$  is quasi-stable if and only if it has no trivial component.*

*Proof.* It is immediate from the definition and the fact that all proper algebraic subgroup of  $G_m$  are finite.  $\square$

Recall that there are two canonical divisors  $D_+$  and  $D_-$  on the  $\mathbf{P}^1$ -bundle  $\Delta$  over  $D$ , we let  $\Delta^+ = \Delta \setminus D_-$  and  $\Delta^- = \Delta \setminus D_+$  and  $q_+ : \Delta^+ \rightarrow D_+$  and  $q_- : \Delta^- \rightarrow D_-$  be the corresponding projections. For any coherent sheaf  $\mathcal{E}$  on  $\Delta$ , we let  $\mathcal{E}^+ = \mathcal{E} \otimes \mathcal{O}_{D_+}$  and  $\mathcal{E}^- = \mathcal{E} \otimes \mathcal{O}_{D_-}$ .

Let  $P \in D_+$  and  $V_P$  be a formal neighborhood of  $P$ . Let  $y = 0$  be the local defining equation of  $D_+$  in  $\Delta^+$ . Suppose  $V_P = \text{Spec } k[[x_i]]$ , then  $q_+^{-1}(V_P) = \text{Spec } k[[x_i]][y]$ . Let  $\mathcal{I}$  be an ideal sheaf on  $\Delta$ . Then the restriction of  $\mathcal{I}$  to  $q_+^{-1}(V_P)$  is an ideal  $I \subset k[[x_i]][y]$ .

**Lemma 2.24.** *Let  $\mathcal{I}$  be an ideal sheaf on  $X[n]_0$  which is perfect along  $D_+ \subset \Delta_k$ . Then  $\Delta_k$  is a trivial component of  $\mathcal{I}$  if and only if  $\mathcal{I} \otimes \mathcal{O}_{\Delta_k^+} = q_+^*(\mathcal{I}_k^+)$ .*

*Proof.* If  $\Delta_k$  is a trivial component of  $\mathcal{I}$ , then every automorphism of  $X[n]_0$  which is a dilation of  $\Delta_k$  and fixes other components must be a categorical isomorphism of  $\mathcal{I}$ . Now we fix a point  $P \in D_+$  and let  $I \subset k[[x_i]][y]$  be the restriction of  $\mathcal{I}$  to  $q_+^{-1}(V_P)$ . Suppose  $I = (f_1(y, x_i), \dots, f_n(y, x_i))$ , let

$$f_1(y, x_i) = g_0(x_i) + yg_1(x_i) + \dots + y^m g_m(x_i).$$

We will show that all  $g_k(x_i)$  are contained in  $I$ . Since  $f_1(\lambda y, x_i) \in I$  for any scalar  $\lambda \neq 0$ ,  $f_1(\lambda y, x_i) - f_1(y, x_i)$  is also contained in  $I$ , that is

$$y(\lambda - 1)g_1(x_i) + y^2(\lambda^2 - 1)g_2(x_i) + \dots + y^m(\lambda^m - 1)g_m(x_i) \in I.$$

Since  $I$  is perfect along  $D_+$ , we have

$$(\lambda - 1)g_1(x_i) + y(\lambda^2 - 1)g_2(x_i) + \dots + y^{m-1}(\lambda^m - 1)g_m(x_i) \in I.$$

By the infinitely many choices of  $\lambda$ , we can continue this process and obtain that  $g_m(x_i) \in I$ . Therefore, all  $g_i(x_i)$  are contained in  $I$ . It implies that  $I$  is generated by formal series of  $x_i$ . Since this is true for every point  $P \in D_+$ , we get  $I = q_+^*(I^+)$  with  $I^+ = I|_{D_+}$ .

Conversely, if  $\mathcal{I} \otimes \mathcal{O}_{\Delta_k^+} = q_+^*(\mathcal{I}_k^+)$ , then at every point  $P \in D_+$ ,  $I = q_+^*(I^+)$ . It implies that  $I = (f_1(x_i), \dots, f_n(x_i))$ . It is obvious that  $I$  is invariant by every dilation of  $\Delta_k$ . Hence  $\Delta_k$  is a trivial component.  $\square$

**Lemma 2.25.** *Fix a homology class  $\Gamma \in H_*(X, \mathbf{Z})$ . There exists an integer  $N$ , such that if  $n > N$ , then there is no quasi-stable ideal sheaf on  $X[n]_0$  of type  $\Gamma$ .*

*Proof.* For the purpose of Donaldson-Thomas invariants, we prove this lemma in the case of ideal sheaves of one dimensional subschemes in threefolds.

By Riemann-Roch theorem, to fix all Chern classes  $c_i(\mathcal{I})$ , it is equivalent to fix  $c_1(\mathcal{I})$ ,  $c_2(\mathcal{I})$  and  $\chi(\mathcal{I})$ . Since  $\chi(\mathcal{O}_\Delta) = \chi(\mathcal{O}_D)$ , we have  $\chi(\mathcal{O}_{X[n]_0}) = \chi(\mathcal{O}_{X_0})$  for every  $n \geq 0$ . Let  $Z$  be the closed subscheme defined by  $\mathcal{I}$ . Then the Chern classes  $c_i(\mathcal{O}_Z)$  and  $\chi(\mathcal{O}_Z)$  are all fixed.

Now we turn to the case of threefolds. Let  $\mathcal{I}$  be an ideal sheaf of a one dimensional closed subscheme  $Z \subset X[n]_0$ . Then  $c_1(\mathcal{O}_Z) = 0$ ,  $c_2(\mathcal{O}_Z) = -[Z]$ . Let  $Z_k = Z \cap \Delta_k$  and  $Z_{D_k} = Z \cap D_k$ .

Next we show the following fact:  $\Delta_k$  is not a trivial component of  $\mathcal{I}$  if and only if  $Z_k$  falls into one of the following cases:

- (1)  $[Z_k] \neq mF$  for any  $m$ , where  $F$  is the fibre class of  $\Delta_k$ .
- (2)  $[Z_k] = mF$  for some  $m$  and  $Z_k$  has zero dimensional torsion part which does not support on distinguished divisors.
- (3)  $[Z_k] = mF$  for some  $m$  and  $\chi(\mathcal{O}_{Z_k^p}) > m$ , where  $Z_k^p$  is the maximal pure subscheme of  $Z_k$ .

The first two cases are immediate from the definition. For case (3), we may assume without loss of generality that  $Z_k$  is irreducible and pure so that  $Z_k^p = Z_k$ . For simplicity, we assume  $m = 2$ . The general case can be proved similarly and by induction on  $m$ . Let  $Z_k^r$  be the reduced part of  $Z_k$ . Then  $Z_k^r \cong F = \mathbf{P}^1$  and there is a canonical exact sequence

$$0 \longrightarrow \mathcal{J} \longrightarrow \mathcal{O}_{Z_k} \longrightarrow \mathcal{O}_{Z_k^r} \longrightarrow 0,$$

where  $\mathcal{J}$  is the ideal sheaf of  $Z_k^r$  in  $Z_k$  with  $\mathcal{J}^2 = 0$ .  $\Delta_k$  is a trivial component of  $Z_k$  if and only if  $\mathcal{J}$  and  $Z_k^r$  are invariant under all dilation  $\lambda$  of  $\Delta_k$ . Let  $\tilde{\mathcal{J}}$  be the ideal sheaf of  $Z_k^r$  in  $\Delta$ . Then there is a surjective homomorphism  $\tilde{\mathcal{J}} \rightarrow \mathcal{J}$ . It induces a nonzero homomorphism  $\tilde{\mathcal{J}}/\tilde{\mathcal{J}}^2 \rightarrow \mathcal{J}$  because  $\mathcal{J}^2 = 0$ . Notice that  $\tilde{\mathcal{J}}/\tilde{\mathcal{J}}^2$  is the conormal bundle of  $Z_k^r$  in  $\Delta$ , it is a direct sum of trivial bundles. Since  $\mathcal{J}$  is a bundle over  $Z_k^r$  and the existence of a nonzero homomorphism  $\tilde{\mathcal{J}}/\tilde{\mathcal{J}}^2 \rightarrow \mathcal{J}$ , the degree of  $\mathcal{J}$  must be nonnegative. Now if  $\Delta_k$  is not a trivial component, the ideal sheaf  $\mathcal{J}$  is not invariant by all dilations. Hence the degree of  $\mathcal{J}$  is positive. Therefore

$$\chi(\mathcal{O}_{Z_k}) = \chi(\mathcal{O}_{Z_k^r}) + \chi(\mathcal{J}) = \chi(\mathbf{P}^1) + \chi(\mathbf{P}^1) + \deg \mathcal{J} > 2.$$

If  $\mathcal{I}$  is a quasi-stable ideal sheaf on  $X[n]_0$ , then  $\mathcal{I}$  has no trivial component. Now we show that there is an upper bound of  $n$ . Since  $\sum_{k=0}^{n+1} p_*[Z_k] = \Gamma_2$  and every class  $p_*[Z_k]$  is positive or zero, we see that there is an integer  $M$  such that the number of components  $\Delta_k$  such that  $p_*[Z_k] \neq 0$  is not greater than  $M$ . A class  $p_*[Z_k]$  is zero if and only if  $[Z_k] = mF$  for some integer  $m > 0$ . Next we show that the number of components  $\Delta_k$  with  $[Z_k] = mF$  is also bounded from above. Since for  $[Z_k] = mF$  (case (2) and (3)), we always have  $\chi(\mathcal{O}_{Z_k}) > \chi(\mathcal{O}_{Z_{D_k}})$ , by  $\chi(\mathcal{O}_Z) = \sum \chi(\mathcal{O}_{Z_k}) - \sum \chi(\mathcal{O}_{Z_{D_k}})$ , we know the number of nontrivial components  $\Delta_k$  with  $[Z_k] = mF$  is bounded.

Combine the discussion, we see that after fixing the topological type  $\Gamma$ , there is an upper bound  $N$  such that if  $n > N$ , then there is no quasi-stable ideal sheaf on  $X[n]_0$ . □

**2.5. The stack of stable perfect ideal sheaves.** First we define the moduli stack  $\mathfrak{I}_{X/C}^\Gamma$  of stable perfect ideal sheaves of type  $\beta$ . An object in  $\mathfrak{I}_{X/C}^\Gamma$  is a pair  $(\mathcal{X}/S, \mathcal{I})$  which consists of an expanded degeneration  $\mathcal{X}$  over a  $C$ -scheme  $S$  and a family of stable perfect ideal sheaves  $\mathcal{I}$  of type  $\beta$  on  $\mathcal{X}/S$ . Let  $\xi_1 = (\mathcal{X}_1/S_1, \mathcal{I}_1)$  and  $\xi_2 = (\mathcal{X}_2/S_2, \mathcal{I}_2)$  be two objects in  $\mathfrak{I}_{X/C}^\Gamma$ . An arrow  $r : \xi_1 \rightarrow \xi_2$  is defined to be a pair  $(u, \psi)$  of which  $u : \mathcal{X}_1/S_1 \rightarrow \mathcal{X}_2/S_2$  is an arrow in the stack  $\mathfrak{X}$  of expanded

degeneration and  $\psi : \mathcal{I}_1 \rightarrow u^*\mathcal{I}_2$  is an isomorphism of ideal sheaves on  $\mathcal{X}_1/S_1$ . There is a forgetful functor  $F : \mathfrak{J}_{X/C}^\Gamma \rightarrow \text{Sch}/C$  which sends  $(\mathcal{X}/S, \mathcal{I})$  to the base scheme  $S$ . It is straightforward to verify that  $\mathfrak{J}_{X/C}^\Gamma$  is a groupoid over  $C$ . In the sequel, We use  $\mathfrak{J}_{X/C}^\Gamma(S)$  to denote the subgroupoid of stable perfect ideal sheaves on  $\mathfrak{X}(S)$ . In particular,  $\mathfrak{J}_{X/C}^\Gamma(\mathbf{0})$  is the subgroupoid of stable perfect ideal sheaves on the stack  $\mathfrak{X}_0$  of semistable models of  $X_0$ .

**Proposition 2.26.**  $\mathfrak{J}_{X/C}^\Gamma$  is a Deligne-Mumford stack of finite type over  $C$ .

*Proof.* First we show that  $\mathfrak{J}_{X/C}^\Gamma$  is a stack. By the definition of stacks, we need to verify the following conditions([13]):

(1) For any scheme  $S$  in  $\text{Sch}/C$  and two families  $\xi_1, \xi_2$  of stable perfect ideal sheaves in  $\mathfrak{J}_{X/C}^\Gamma(S)$ , the functor

$$\text{Isom}_S(\xi_1, \xi_2) : \text{Sch}/C \rightarrow (\text{Sets}),$$

which associates to any morphism  $\phi : T \rightarrow S$  the set of isomorphisms in  $\mathfrak{J}_{X/C}^\Gamma(T)$  between  $\phi^*\xi_1$  and  $\phi^*\xi_2$  is a sheaf in the étale topology.

(2) Let  $\{S_i \rightarrow S\}$  be a covering of  $S \in \text{Sch}/C$  in the étale topology. Let  $\xi_i \in \mathfrak{J}_{X/C}^\Gamma(S_i)$  and let  $\varphi_{ij} : \xi_j|_{S_i \times_S S_j} \rightarrow \xi_i|_{S_i \times_S S_j}$  be isomorphisms in  $\mathfrak{J}_{X/C}^\Gamma(S_i \times_S S_j)$  satisfying the cocycle condition. Then there is  $\xi \in \mathfrak{J}_{X/C}^\Gamma(S)$  with isomorphism  $\psi_i : \xi|_{S_i} \rightarrow \xi_i$  such that

$$\varphi_{ij} = (\psi_i|_{S_i \times_S S_j}) \circ (\psi_j|_{S_i \times_S S_j})^{-1}.$$

Since stable perfect ideal sheaves have finite automorphism groups, by the result of Grothendieck [4],  $\text{Isom}_S(\xi_1, \xi_2)$  is represented by a finite group scheme. The proof of (1) and (2) follows from the standard argument in descent theory.

To prove  $\mathfrak{J}_{X/C}^\Gamma$  is algebraic, it suffices to show that  $\mathfrak{J}_{X/C}^\Gamma$  admits an étale cover by a scheme of finite type.

Let  $\mathbf{Hilb}_{X[n]/C[n]}$  be the Hilbert scheme of one dimensional subschemes  $Z$  such that  $\mathcal{I}_Z$  has topological type  $\Gamma$ . Let  $\mathbf{Hilb}_{X[n]/C[n]}^p$  be the open subscheme of  $\mathbf{Hilb}_{X[n]/C[n]}$  parameterizing stable perfect ideal sheaves  $\mathcal{I}_Z$ . Then there is a natural morphism

$$\Psi_n : \mathbf{Hilb}_{X[n]/C[n]}^p \longrightarrow \mathfrak{J}_{X/C}^\Gamma.$$

It is obvious that  $\Psi_n$  is an open embedding.

From Lemma 2.25, there is an integer  $N$  so that there is no stable perfect ideal sheaf on  $X[n]_0$  for  $n > N$ , we get a surjective morphism

$$\bigsqcup_{n=0}^N \mathbf{Hilb}_{X[n]/C[n]}^p \longrightarrow \mathfrak{J}_{X/C}^\Gamma.$$

For any closed point  $p \in \mathfrak{J}_{X/C}^\Gamma$ , there exists an  $n$  such that  $p$  is contained in the image of  $\Psi_n$ .  $\mathbf{Hilb}_{X[n]/C[n]}^p$  admits a natural  $G[n]$  action, by the definition of stability, the stabilizer of this action is finite and reduced. Let  $p'$  be a point in the inverse image of  $p$ . Then there is a slice  $S \subset \mathbf{Hilb}_{X[n]/C[n]}^p$  containing  $p'$ , such that the composition  $S \rightarrow \mathbf{Hilb}_{X[n]/C[n]}^p \rightarrow \mathfrak{J}_{X/C}^\Gamma$  is étale. Finally, since  $\mathfrak{J}_{X/C}^\Gamma$  can be covered by finitely many such schemes. We let  $U$  be the finite union of these schemes. Then there is a surjective étale morphism  $U \rightarrow \mathfrak{J}_{X/C}^\Gamma$ .  $\square$



Now we recall the valuative criterion of separatedness and properness of a morphism between two stacks [2].

**Proposition 2.27** (Valuative criterion for separatedness). *Let  $f : \mathfrak{C}_1 \rightarrow \mathfrak{C}_2$  be a morphism between two stacks. Then  $f$  is separated if and only if, for any complete discrete valuation ring  $R$  with algebraically closed residue field and any commutative diagram*

$$\begin{array}{ccc} & & \mathfrak{C}_1 \\ g_1 \nearrow & & \downarrow f \\ \text{Spec } R & \longrightarrow & \mathfrak{C}_2 \\ g_2 \nearrow & & \end{array}$$

*any isomorphism between the restrictions of  $g_1$  and  $g_2$  to the generic point of  $\text{Spec } R$  can be extended to an isomorphism between  $g_1$  and  $g_2$ .*

**Proposition 2.28** (Valuative criterion for properness). *If  $f$  is separated then  $f$  is proper if and only if, for any discrete valuation ring  $R$  with field of fraction  $K$  and any commutative diagram*

$$\begin{array}{ccc} \text{Spec } K & \xrightarrow{g} & \mathfrak{C}_1 \\ \downarrow & & \downarrow f \\ \text{Spec } R & \longrightarrow & \mathfrak{C}_2 \end{array}$$

*there exists a finite extension  $K'$  of  $K$  such that  $g$  extends to  $\text{Spec } R'$ , where  $R'$  is the integral closure of  $R$  in  $K'$ .*

$$\begin{array}{ccccc} \text{Spec } K' & \longrightarrow & \text{Spec } K & \xrightarrow{g} & \mathfrak{C}_1 \\ \downarrow & & \downarrow & \nearrow & \downarrow f \\ \text{Spec } R' & \longrightarrow & \text{Spec } R & \longrightarrow & \mathfrak{C}_2 \end{array}$$

In the remainder of this section, we will show that  $\mathfrak{I}_{X/C}^\Gamma$  is proper over  $C$ .

Let us begin by recalling some notation. Let  $S$  be a one dimensional regular scheme over  $C$  with a closed point  $\eta_0$  lying over  $\mathbf{0} \in C$ . Let  $\mathcal{X}$  be the effective degeneration over  $S$  associated to a morphism  $f : S \rightarrow C[n]$  with  $f(\eta_0) = \mathbf{0} \in C[n]$ . Let  $\iota_{n+1} : C[n] \rightarrow C[n+1]$  be a standard embedding. Let  $S^* = S - \eta_0$  and  $\rho : S^* \rightarrow G[n+1]$  be a morphism. Then there is an effective degeneration over  $S^*$  associated to the morphism  $f_\rho = (\iota_{n+1} \circ f)^\rho : S^* \rightarrow C[n+1]$ . We choose  $\rho$  appropriately such that  $f_\rho$  can be extended to  $S$  and  $f_\rho(\eta_0) = \mathbf{0} \in C[n+1]$ . We denote by  $\mathcal{X}^\rho$  the resulting effective degeneration over  $S$ . The fiber of  $\mathcal{X}^\rho$  over  $\eta_0$  is a semistable model of  $X_0$  with length  $n+1$ . There is a morphism  $\mathcal{X}^\rho/S \rightarrow \mathcal{X}/S$  whose restriction to  $S^* \subset S$  is an isomorphism. Now we let  $\mathcal{I}$  be a family of ideal sheaves on  $\mathcal{X}/S$ . By the isomorphism  $\mathcal{X}^\rho \times_S S^* \rightarrow \mathcal{X} \times_S S^*$ , we obtain a corresponding family of ideal sheaves on  $\mathcal{X}^\rho \times_S S^*$ . By the properness of the Hilbert scheme, this family extends uniquely to a family of ideal sheaves on  $\mathcal{X}^\rho/S$ , which is denoted by  $\mathcal{I}^\rho$ .

The following is a key technical lemma in the proof of properness. The proof of this lemma will be given in the last section of this paper.

**Lemma 2.29.** *Keep the notation as above. Let  $\mathcal{I}$  be a family of ideal sheaves on  $\mathcal{X}/S$  such that the restriction of  $\mathcal{I}$  to  $\mathcal{X} \times_S S^*$  is a family of perfect ideal sheaves.*

Let  $\mathcal{I}_0 = \mathcal{I} \otimes_{\mathcal{O}_S} k_{\eta_0}$  be an ideal sheaf on  $X[n]_0$ . We assume that  $\mathcal{I}_0$  is not perfect along  $D_k$  for some  $k$ . Then there is a finite morphism  $(S_1, \eta'_0) \rightarrow (S, \eta_0)$  totally ramified at  $\eta_0$ , and a morphism  $\rho_r : S_1^* \rightarrow G[n+1]$ , so that  $f_\rho : S_1^* \rightarrow C[n+1]$  can be extended to  $S_1$  such that  $f_\rho(\eta'_0) = \mathbf{0}$ , and such that the resulting family  $\mathcal{I}^\rho$  on  $\mathcal{X}^\rho/S$  satisfies the following properties:

- (1) The ideal sheaf  $\mathcal{I}_0^\rho$  is perfect along  $D_k \subset X[n+1]_0$ .
- (2)  $\Delta_k$  is not a trivial component of  $\mathcal{I}_0^\rho$ .

In the following application of this lemma, we let  $S = \text{Spec } R$  and  $S^* = \text{Spec } K$ , where  $R$  is a discrete valuation ring and  $K$  is the field of fractions of  $R$ .

**Proposition 2.30.** *The stack  $\mathfrak{J}_{X/C}^\Gamma$  is separated over  $C$ .*

*Proof.* Let  $R$  be a complete discrete valuation ring and  $S = \text{Spec } R$ . Let  $g_1, g_2 : S \rightarrow \mathfrak{J}_{X/C}^\Gamma$  be two morphisms. Because of the canonical equivalence between the groupoid  $\text{Hom}(S, \mathfrak{J}_{X/C}^\Gamma)$  of morphisms and the fiber groupoid  $\mathfrak{J}_{X/C}^\Gamma(S)$ , there are two objects  $\xi_1, \xi_2$  in  $\mathfrak{J}_{X/C}^\Gamma(S)$  corresponding to the morphisms  $g_1, g_2$  respectively. Let  $\eta$  be the generic point of  $S$ . Let  $\xi_{i,\eta}$  be the restriction of the object  $\xi_i$  to  $\eta$ , i.e.  $\xi_{i,\eta} = \xi_i \times_S \eta$ . By the valuative criterion, to prove that  $\mathfrak{J}_{X/C}^\Gamma$  is separated over  $C$ , it suffices to show that every isomorphism  $r_\eta : \xi_{1,\eta} \rightarrow \xi_{2,\eta}$  can be extended to an isomorphism  $r : \xi_1 \rightarrow \xi_2$ .

Since  $\mathcal{X}_{1,\eta} \cong \mathcal{X}_{2,\eta}$ , we denote this isomorphism class of expanded degeneration over  $\text{Spec } K$  by  $\mathcal{X}_\eta$ . According to the smoothness of  $\mathcal{X}_\eta$ , the argument is slightly different. Next we discuss the case  $\mathcal{X}_\eta$  is smooth.

We further assume that  $\mathcal{X}_i$  is associated to a morphism  $f_i : S \rightarrow C[n_i]$ . Without loss of generality, let  $n_1 \leq n_2$ . We also assume  $n_2$  is minimal among all possible choices. We assume that  $\xi_i$  is represented by a family of ideal sheaves  $\mathcal{I}_i$  on  $\mathcal{X}_i/S$ . Let  $r_\eta : \xi_{1,\eta} \rightarrow \xi_{2,\eta}$  be an isomorphism. let  $\iota : C[n_1] \rightarrow C[n_2]$  be a standard embedding. By Lemma 1.2,  $r_\eta$  is induced by a  $G[n_2]$  action on  $C[n_2]$  via a morphism  $\rho_\eta : \text{Spec } K \rightarrow G[n_2]$ , that is,  $\iota \circ f_1(\eta) = f_2^\rho(\eta)$ . Let  $\tilde{f}_i = p_i \circ f_i : S \rightarrow C$  where  $p_i : C[n_i] \rightarrow C$  is the canonical projection. Since  $\tilde{f}_1(\eta) = \tilde{f}_2(\eta)$  and  $C$  is separated, we get  $\tilde{f}_1(\eta_0) = \tilde{f}_2(\eta_0)$ . It implies that if one of  $\mathcal{X}_{i,\eta_0}$  is smooth, then both of them are smooth. Suppose the fibers  $\mathcal{X}_{i,\eta_0}$  are smooth, Then  $\iota \circ f_1(S)$  and  $f_2(S)$  are both contained in  $C[n_2]^*$ . Since  $C[n_2]^*$  is a principal fiber bundle over  $C^*$  and  $\tilde{f}_1(S) = \tilde{f}_2(S)$ , there is a morphism  $\rho : S \rightarrow G[n_2]$  extending  $\rho_\eta$ , that is,  $\iota \circ f_1 = f_2^\rho$ . It implies that  $r_\eta : \xi_{1,\eta} \rightarrow \xi_{2,\eta}$  can be extended to an isomorphism  $r : \xi_1 \rightarrow \xi_2$ .

Next we assume that both  $\mathcal{X}_{1,\eta_0}$  and  $\mathcal{X}_{2,\eta_0}$  are singular. We first deal with the case  $f_1(\eta_0) = f_2(\eta_0) = \mathbf{0} \in C[n_2]$ . Since the isomorphism  $r_\eta$  is induced from a  $G[n_2]$  action via a morphism  $\rho_\eta : \text{Spec } K \rightarrow G[n_2]$ , we choose  $s$  as a local uniformizer of  $R$ , then  $\rho_\eta$  can be defined by

$$\rho_\eta^*(\sigma_1) = c_1 s^{a_1}, \quad \dots, \quad \rho_\eta^*(\sigma_{n_2}) = c_{n_2} s^{a_{n_2}},$$

where  $c_i$  are units in  $R$  and  $a_i$  are integers.  $\rho_\eta$  can be extended to  $S$  if and only if all  $a_i = 0$ .

Now we show that  $a_1 = 0$ . Let  $\iota_1 : C[n_2 - 1] \rightarrow C[n_2]$  be a standard embedding associated to  $I = \{2, \dots, n_2 + 1\}$ . We can choose a morphism  $g : S \rightarrow C[n_2 - 1]$  such that the effective degeneration  $\iota_1 \circ g : \text{Spec } K \rightarrow C[n_2]$  is equivalent to  $f_2 : \text{Spec } K \rightarrow C[n_2]$ . We claim that the family is not perfect along  $\Delta_k$ . For otherwise, by the lemma, we know that the original family has a trivial component  $\Delta_k$ , which

contradicts to the stability of  $\mathcal{I}_0$ . From Lemma 2.29, there is a unique morphism  $\rho_a : \text{Spec } K \rightarrow G[n_2]$ , such that  $\mathcal{I}^{\rho_a}$  is a family of stable perfect ideal sheaves on  $\mathcal{X}_2$ . Since  $\rho = \rho_a \rho_\eta$  also gives rise to a family of stable perfect ideal sheaves, by the uniqueness in Lemma 2.29, we obtain  $a + a_1 = a$  and hence  $a_1 = 0$ . Likewise, we have  $a_i = 0$  for all  $i > 1$ .

Finally, we assume both  $f_i(\eta_0) \neq \mathbf{0}$  and derive a contradiction. The proof of the case in which only one of  $f_i(\eta_0) \neq \mathbf{0}$  is similar. It is clear that we can assume  $f_i(\eta_0)$  lay on different coordinate hyperplanes of  $C[n_2]$ , for otherwise, we can use projection  $C[n_2] \rightarrow C[n_2 - 1]$  to obtain new isomorphic families which contradicts to the minimality of  $n_2$ . Using the same argument as above, there are morphisms  $\rho_1 : \text{Spec } K \rightarrow G[n_2]$  and  $\rho_2 : \text{Spec } K \rightarrow G[n_2]$  such that  $f_1^{\rho_1}(\eta) = f_2^{\rho_2}(\eta)$ . Now  $\mathcal{I}_1^{\rho_1}$  and  $\mathcal{I}_2^{\rho_2}$  are families on the same expanded degeneration  $\mathcal{X}_2/S$ . The properness of Hilbert scheme implies that  $\mathcal{I}_1^{\rho_1} \otimes k_{\eta_0} \cong \mathcal{I}_2^{\rho_2} \otimes k_{\eta_0}$ . This is impossible since  $\mathcal{I}_1^{\rho_1} \otimes k_{\eta_0}$  and  $\mathcal{I}_2^{\rho_2} \otimes k_{\eta_0}$  have different trivial components.  $\square$

**Proposition 2.31.** *The stack  $\mathfrak{J}_{X/C}^\Gamma$  is proper over  $C$ .*

*Proof.* Let  $R$  be a discrete valuation ring with fraction field  $K$ . Let  $g : \text{Spec } K \rightarrow \mathfrak{J}_{X/C}^\Gamma$  be a morphism. By the canonical equivalence between  $\text{Hom}(\text{Spec } K, \mathfrak{J}_{X/C}^\Gamma)$  and  $\mathfrak{J}_{X/C}^\Gamma(\text{Spec } K)$ , we get an object  $\xi_\eta = (\mathcal{X}_\eta / \text{Spec } K, \mathcal{I})$  in  $\mathfrak{J}_{X/C}^\Gamma$  corresponding to the morphism  $g : \text{Spec } K \rightarrow \mathfrak{J}_{X/C}^\Gamma$ .

Given a commutative diagram

$$\begin{array}{ccc} \text{Spec } K & \xrightarrow{g} & \mathfrak{J}_{X/C}^\Gamma \\ \downarrow & & \downarrow F \\ \text{Spec } R & \longrightarrow & C, \end{array}$$

according to the valuation criterion of properness, we need to show that probably after a base change  $q : \text{Spec } K' \rightarrow \text{Spec } K$ , the family of stable perfect ideal sheaves  $q^*\mathcal{I}$  on  $\mathcal{X}_\eta \times_{\text{Spec } K} \text{Spec } K'$  parameterized by  $\text{Spec } K'$  can be extended to  $\text{Spec } R'$ , where  $R'$  is the integral closure of  $R$  in  $K'$ .

Assume that the expanded degeneration  $\mathcal{X}_\eta$  is associated to a morphism  $f : \text{Spec } K \rightarrow C[n]$  for some  $n$ . Let  $p : C[n] \rightarrow C$  be the canonical projection. we get a composition  $\tilde{f} = p \circ f : \text{Spec } K \rightarrow C$ . The above commutative diagram says that  $\tilde{f}$  can be extended to  $S = \text{Spec } R$ . We now separate the following discussion into two cases.

**Case 1.**  $\tilde{f} : S \rightarrow C$  does not factor through  $\mathbf{0} \in C$ .

In this case, the generic fiber  $\mathcal{X}_\eta$  is smooth and  $\tilde{f}$  induces an associated expanded degeneration  $\mathcal{X}'$  over  $S$ , such that there is an isomorphism  $\psi : \mathcal{X}' \times_S \text{Spec } K \cong \mathcal{X}_\eta$ . Now since ideal sheaves are stable in the ordinary sense, the properness of the moduli scheme of ordinary stable sheaves implies that the family  $\psi^*\mathcal{I}$  can be extended to a family of ideal sheaves  $\tilde{\mathcal{I}}$  on  $\mathcal{X}'$ . Next we need to check the perfectness of the ideal sheaf  $\tilde{\mathcal{I}}_0$  on  $\mathcal{X}'_{\eta_0}$ . If  $\mathcal{X}'_{\eta_0}$  is smooth, all ideal sheaves on  $\mathcal{X}'_{\eta_0}$  are automatically perfect which completes the proof. Next, if  $\mathcal{X}'_{\eta_0}$  is not smooth, then  $\mathcal{X}'_{\eta_0} \cong X[m]_0$  for some  $m$ . We can let  $\mathcal{X}'$  be represented by a morphism  $S \rightarrow C[m]$ . By the properness of Hilbert schemes,  $\mathcal{I}$  can be extended to a family of ideal sheaves on  $\mathcal{X}'$ . We say  $\mathcal{I}$  admits a partial extension to  $\mathcal{X}'$ , if  $\mathcal{I}$  extends to a family of quasi-stable (not necessarily perfect) ideal sheaves over  $S$ . By Lemma

2.25, there is a constant  $N$  such that  $n \leq N$ . Hence the set of integers  $n$  such that  $\mathcal{I}$  admits a partial extension to  $\mathcal{X}'$  is bounded from above. Let  $n_0$  be the largest integer in this finite set which corresponds to an expanded degeneration  $\mathcal{X}'$  given by a morphism  $S \rightarrow C[n_0]$ , and a family of ideal sheaves  $\mathcal{I}'$  on  $\mathcal{X}'$  such that  $\mathcal{I}'_0$  is quasi-stable. We claim that  $\mathcal{I}'_0$  is perfect and thus stable because it is already quasi-stable. Now we assume  $\mathcal{I}'_0$  is not perfect and derive a contradiction. Assume further  $\mathcal{I}'_0$  is not perfect along  $D_k$  for some  $k$ . Now by Lemma 2.29, there exists a base change  $S' \rightarrow S$  and a morphism  $S' \rightarrow C[n_0 + 1]$ , such that the ideal sheaf  $\mathcal{I}$  is perfect along  $D_k$  and  $\mathcal{I}$  has no trivial component. Clearly it contradicts to the maximality of  $n_0$ . Hence  $\mathcal{I}$  is already perfect.

**Case 2.**  $\tilde{f} : \text{Spec } R \rightarrow C$  factors through  $\mathbf{0} \in C$ .

In this case,  $f : \text{Spec } K \rightarrow C[n]$  must send  $\text{Spec } K$  into a coordinate hyperplane of  $C[n]$ . It implies that  $\mathcal{X}$  is a constant family over  $\text{Spec } K$ , i.e.  $\mathcal{X} = X[m]_0 \times \text{Spec } K$  for some  $m$ . There is a trivial extension of this family to  $\text{Spec } R$ , which is denoted by  $\mathcal{X}'$ . For this constant family  $\mathcal{X}'$ , we can always find an associated morphism  $f' : S \rightarrow C[n]$  which is not a constant morphism. Now we can follow a similar argument as in case 1 and use a variation of lemma 2.29 to obtain the result.  $\square$

Combine Proposition 2.26, 2.30 and 2.31, we obtain the following

**Theorem 2.32.** *The moduli stack  $\mathfrak{I}_{X/C}^\Gamma$  is a separated and proper Deligne-Mumford stack of finite type over  $C$ .*

### 3. THE STACK OF STABLE RELATIVE IDEAL SHEAVES

In this section, we first recall the notion of extended pairs and the decomposition of semistable model  $X[n]_0$  into extended pairs. Next we define relative ideal sheaves on extended pairs, and the stack of stable relative ideal sheaves which has similar properties as the stack of stable perfect ideal sheaves studied in the previous section.

**3.1. Expanded pairs.** Let  $(Y, D)$  be a pair of a smooth variety  $Y$  and a smooth divisor  $D \subset Y$ . Let  $N = N_{D/Y}$  be the normal bundle of  $D$  in  $Y$ . We now define the extended pairs of  $(Y, D)$ . To begin with, recall that  $\Delta = \mathbf{P}(N \oplus \mathbf{1}_D)$  is a  $\mathbf{P}^1$  bundle over  $D$  with two canonical divisors  $D_+$  and  $D_-$  satisfying that  $N_{D_+/\Delta} \cong N$  and  $N_{D_-/\Delta} \cong N^\vee$ . Again we use  $\Delta_i$  to denote the  $i^{\text{th}}$  copy of  $\Delta$  for  $i > 0$ . An extended pair with length  $n$ , denoted as  $(Y[n]_0, D[n]_0)$ , consists of a scheme  $Y[n]_0$  which is a chain of smooth irreducible components  $Y, \Delta_1, \dots, \Delta_n$  intersecting transversally, i.e.

$$Y[n]_0 = Y \cup \Delta_1 \cup \dots \cup \Delta_n,$$

and a divisor  $D[n]_0 = D_+$  on the last component  $\Delta_n$ . For convenience, we often denote the irreducible component  $Y \subset Y[n]_0$  by  $\Delta_0$  and let  $D_i = \Delta_{i-1} \cap \Delta_i$  for  $i > 0$ . The union  $\Delta_1 \cup \dots \cup \Delta_n$  is usually called the tail of  $Y[n]_0$ . Since  $\Delta$  is a projective bundle over  $D$ , there is a canonical surjective morphism from the tail of  $Y[n]_0$  to  $D$ . It therefore implies that there is a canonical projection  $p_n : Y[n]_0 \rightarrow Y$ , which is an identity morphism on the component  $Y$ , and sends the tail of  $Y[n]_0$  to the distinguished divisor  $D \subset Y$ .

Let  $\xi_1 = (Y[n]_0, D[n]_0)$  and  $\xi_2 = (Y[m]_0, D[m]_0)$  be two extended pairs. A morphism  $\xi_1 \rightarrow \xi_2$  is a morphism  $u : Y[n]_0 \rightarrow Y[m]_0$  such that the image of  $D[n]_0$  is  $D[m]_0$ , which is compatible with their canonical projections, i.e.  $p_m \circ u = p_n$ .

Because the group of automorphisms of  $\Delta$  which fix  $D_+$  and  $D_-$  is  $G_m$ , it is easy to verify that the automorphism group of an extended pair  $(Y[n]_0, D[n]_0)$  is  $G[n]$ .

Next we are naturally led to families of extended pairs. To begin with, we first describe the construction of relative standard model which is a family  $(Y[n], D[n])$  over affine space  $\mathbf{A}^n$ . As before, let  $(Y, D)$  be a fixed pair. The relative standard model  $(Y[n], D[n])$  of  $(Y, D)$  is defined inductively as follows. For  $n = 0$ ,  $(Y[0], D[0]) = (Y, D)$ . For  $n = 1$ , we let  $\pi_1 : Y[1] \rightarrow Y \times \mathbf{A}^1$  be the blowup of  $Y \times \mathbf{A}^1$  along the codimension 2 subvariety  $D \times \mathbf{0} \subset Y \times \mathbf{A}^1$ . Because the subvariety  $D \times \mathbf{0}$  is isomorphic to  $D$  and its normal bundle in  $Y \times \mathbf{A}^1$  is isomorphic to  $N \oplus \mathbf{1}_D$ , the exceptional divisor of the blowup  $\pi_1$  is isomorphic to the  $\mathbf{P}^1$  bundle  $\Delta$  over  $D$ . Let  $D[1]$  be the proper transformation of  $D \times \mathbf{A}^1$ . Then  $D[1]$  is a smooth divisor in  $Y[1]$ . We then have a morphism  $\pi_1 : (Y[1], D[1]) \rightarrow (Y, D) \times \mathbf{A}^1$ , where  $(Y, D) \times \mathbf{A}^1 = (Y \times \mathbf{A}^1, D \times \mathbf{A}^1)$  is a constant family of  $(Y, D)$ . Let  $q_1 : (Y, D) \times \mathbf{A}^1 \rightarrow (Y, D)$  and  $q_2 : (Y, D) \times \mathbf{A}^1 \rightarrow \mathbf{A}^1$  be the projections. Then  $q_2 \circ \pi_1 : (Y[1], D[1]) \rightarrow \mathbf{A}^1$  is a family of extended pairs parameterized by  $\mathbf{A}^1$ . The fibre of this family over  $t \neq \mathbf{0}$  is isomorphic to  $(Y, D)$ , and the fibre over  $\mathbf{0} \in \mathbf{A}^1$  is an extended pair  $(Y[1]_0, D[1]_0)$ . Suppose we have defined  $(Y[n-1], D[n-1])$ , then  $Y[n]$  is the blowup of  $Y[n-1] \times \mathbf{A}^1$  along the subvariety  $D[n-1] \times \mathbf{0}$  and  $D[n]$  is the proper transformation of  $D[n-1] \times \mathbf{A}^1$ . From this construction, we know that  $(Y[n], D[n])$  is a family of extended pairs over  $\mathbf{A}^n$ , and there is a canonical projection  $(Y[n], D[n]) \rightarrow (Y, D) \times \mathbf{A}^n$ . The fibre of the family  $(Y[n], D[n]) \rightarrow \mathbf{A}^n$  over  $\mathbf{0} \in \mathbf{A}^n$  is exactly the extended pair  $(Y[n]_0, D[n]_0)$ . We also have an isomorphism  $D[n] \cong D \times \mathbf{A}^n$  such that the family  $D[n] \rightarrow \mathbf{A}^n$  coincides with the projection of  $D \times \mathbf{A}^n$  to the second factor. There is an ordering of the coordinate axis  $t_l$  of  $\mathbf{A}^n$ , such that the restriction of  $(Y[n], D[n])$  to  $t_l$ -axis is the smoothing along the divisor  $D_l$  of  $(Y[n]_0, D[n]_0)$ .

An important feature of the relative standard model is that there is a standard  $G[n]$  action on  $(Y[n], D[n])$  which extends the automorphism group  $G[n]$  action on the central fibre  $(Y[n]_0, D[n]_0)$ . This group action can also be defined inductively. Let  $\mathbf{A}^1$  be an affine line with a distinguished point  $\mathbf{0}$ . We let  $\sigma$  be a  $G[1]$  action on  $\mathbf{A}^1$  with  $\mathbf{0}$  fixed. It induces a  $G[1]$  action on the product  $Y \times \mathbf{A}^1$ , which fixes the fiber  $Y \times \mathbf{0}$ . Because  $D \times \mathbf{0} \subset Y \times \mathbf{0}$  is fixed by the group action, and  $Y[1]$  is the blowup of  $Y \times \mathbf{A}^1$  along  $D \times \mathbf{0}$ , the  $G[1]$  action  $Y \times \mathbf{A}^1$  can be lifted to  $Y[1]$ . It is direct to verify that the restriction of the action on the central fiber is the automorphism group action of  $(Y[1]_0, D[1]_0)$ . Similarly, if we have defined a  $G[n-1]$  action on  $(Y[n-1], D[n-1])$ , we get a  $G[n] = G[n-1] \times G[1]$  action on the product  $(Y[n-1], D[n-1]) \times \mathbf{A}^1$  via the  $G[1]$  action on  $\mathbf{A}^1$  mentioned. The lifting of this action to the blowup  $Y[n]$  along  $D[n-1] \times \mathbf{0}$  is the required  $G[n]$  action on  $(Y[n], D[n])$ .

Now we define effective families of extended pairs, which are local models of general families of extended pairs.

**Definition 3.1.** *Let  $S$  be a scheme. An effective family of extended pairs over  $S$  is a family  $(\mathcal{Y}, \mathcal{D}) \rightarrow S$  associated to a morphism  $\tau : S \rightarrow \mathbf{A}^l$  for some  $l > 0$  defined as*

$$\mathcal{Y} = \tau^* Y[l] = Y[l] \times_{\mathbf{A}^l} S, \quad \mathcal{D} = \tau^* D[l] = D[l] \times_{\mathbf{A}^l} S,$$

where  $(Y[l], D[l]) \rightarrow \mathbf{A}^l$  is a relative standard model of  $(Y, D)$ .

Let  $\xi_1 = (\mathcal{Y}_1, \mathcal{D}_1)$  and  $\xi_2 = (\mathcal{Y}_2, \mathcal{D}_2)$  be two effective family of extended pairs over  $S$  which are associated to morphisms  $\tau_1 : S \rightarrow \mathbf{A}^{n_1}$  and  $\tau_2 : S \rightarrow \mathbf{A}^{n_2}$  respectively. Assume  $n_1 \leq n_2$ . An effective arrow  $r : \xi_1 \rightarrow \xi_2$  consists of a standard embedding  $\iota : \mathbf{A}^{n_1} \rightarrow \mathbf{A}^{n_2}$  and a morphism  $\rho : S \rightarrow G[n_2]$ , such that  $(\iota \circ \tau_1)^\rho = \tau_2$ . It is direct to verify that an effective arrow induces an  $S$ -isomorphism of effective pairs compatible to their canonical projections to  $(Y, D) \times S$ . Two effective pairs are equivalent if there is an effective arrow  $\xi_1 \rightarrow \xi_2$  or an effective arrow  $\xi_2 \rightarrow \xi_1$ .

Now we define families of extended pairs.

**Definition 3.2.** *A family of extended pair  $(\mathcal{Y}, \mathcal{D})$  over  $S$  consists of a flat family of schemes  $\mathcal{Y}$  parameterized by  $S$  and a subscheme  $\mathcal{D}$  of  $\mathcal{Y}$ , with a projection  $p : (\mathcal{Y}, \mathcal{D}) \rightarrow (Y, D) \times S$ , satisfying the following condition: There exists an open covering  $\{S_\alpha\}$  of  $S$  such that for each  $\alpha$ , the restriction  $(\mathcal{Y} \times_S S_\alpha, \mathcal{D} \times_S S_\alpha)$  is isomorphic to some effective extended pair over  $S_\alpha$ .*

**Lemma 3.3** ([7]). *Let  $\xi_1 = (\mathcal{Y}_1, \mathcal{D}_1)$  and  $\xi_2 = (\mathcal{Y}_2, \mathcal{D}_2)$  be two extended pairs over  $S$ . Suppose there is an isomorphism  $(\mathcal{Y}_1, \mathcal{D}_1) \cong (\mathcal{Y}_2, \mathcal{D}_2)$ , then for each closed point  $p \in S$ , there is a open neighborhood  $S_0$  of  $p$  and morphisms  $\tau_i : S_0 \rightarrow \mathbf{A}^l$ , such that  $(\mathcal{Y}_i, \mathcal{D}_i) \times_S S_0$  is isomorphic to the associated effective extended pair of  $\tau_i$ , and the induced isomorphism  $(\mathcal{Y}_1, \mathcal{D}_1) \times_S S_0 \cong (\mathcal{Y}_2, \mathcal{D}_2) \times_S S_0$  over  $S_0$  is induced from a sequence of effective arrows between  $\tau_1$  and  $\tau_2$ .*

Next we define the groupoid  $\mathfrak{Y}$  of extended pairs of  $(Y, D)$ . An object in the category  $\mathfrak{Y}$  is a family of extended pair  $(\mathcal{Y}, \mathcal{D})$  over a scheme  $S$ . Let  $\xi_1 = (\mathcal{Y}_1, \mathcal{D}_1)$  and  $\xi_2 = (\mathcal{Y}_2, \mathcal{D}_2)$  be two families of extended pairs over  $S_1$  and  $S_2$  respectively. An arrow  $r : \xi_1 \rightarrow \xi_2$  consists of a morphism  $u : S_1 \rightarrow S_2$  and an isomorphism  $\xi_1 \cong u^* \xi_2$  of families of extended pairs over  $S_1$ . There is a functor  $F : \mathfrak{Y} \rightarrow Sch$  which sends a family  $(\mathcal{Y}, \mathcal{D})$  to its base scheme  $S$ . It is straightforward to verify that the pair of the category  $\mathfrak{Y}$  and the functor  $F$  form a groupoid over  $Sch$ . Furthermore,  $\mathfrak{Y}$  is a stack under the étale topology of  $Sch$ . In the sequel, we use  $\mathfrak{Y}(S)$  to denote the fibre stack over  $S$  which consists of families of extended pairs over  $S$ .

For every family of extended pairs  $(\mathcal{Y}, \mathcal{D})$  over  $S$ , since  $\mathcal{D}$  is a distinguished closed subscheme of  $\mathcal{Y}$  and  $\mathcal{D} \cong D \times S$ , it defines a functor from  $\mathfrak{Y}$  to the stack  $\mathfrak{D}$  associated to the family  $D \rightarrow Spec k$ . Let  $\pi : X \rightarrow C$  be the fixed degeneration mentioned in section 1. There are two pairs  $(Y_1, D_1)$  and  $(Y_2, D_2)$  such that  $X_0 = Y_1 \cup_D Y_2$ . Let  $\mathfrak{Y}_1$  and  $\mathfrak{Y}_2$  be the stack of extended pairs of  $(Y_1, D_1)$  and  $(Y_2, D_2)$  respectively. Since  $D_1 \cong D_2 \cong D$ , we get morphisms  $F_1 : \mathfrak{Y}_1 \rightarrow \mathfrak{D}$  and  $F_2 : \mathfrak{Y}_2 \rightarrow \mathfrak{D}$ . The gluing operation gives a surjective morphism  $G : \mathfrak{Y}_1 \times_{\mathfrak{D}} \mathfrak{Y}_2 \rightarrow \mathfrak{X}_0$ .

**3.2. Stable relative ideal sheaves.** Let  $(Y[n]_0, D[n]_0)$  be an extended pair. We now define relative ideal sheaves on  $(Y[n]_0, D[n]_0)$ .

**Definition 3.4.** *A relative ideal sheaf on  $(Y[n]_0, D[n]_0)$  is a perfect ideal sheaf  $\mathcal{I}$  on  $Y[n]_0$  such that  $\mathcal{I}$  is relative to  $D[n]_0$ .*

**Definition 3.5.** *A relative ideal sheaf  $\mathcal{I}$  on  $(Y[n]_0, D[n]_0)$  is stable, if it is stable as a perfect ideal sheaf.*

The topological type of a relative ideal sheaf on an extended pair can be defined in a similar way. Let  $c(\mathcal{I})$  be the Chern class of a relative ideal sheaf on  $(Y[n]_0, D[n]_0)$  and  $\gamma_i$  be the Poincare dual of the restriction of  $c$  to  $\Delta_i$ . Let  $\gamma = \sum \gamma_i$ . We say  $\mathcal{I}$  has type  $\Gamma$ , denoted as  $\gamma(\mathcal{I}) \sim \Gamma$ , if  $p_*(\gamma) = \Gamma$  with  $p : (Y[n]_0, D[n]_0) \rightarrow (Y, D)$  the canonical projection.

**Definition 3.6.** Fix a class  $\Gamma \in H_*(Y, \mathbf{Z})$ . Let  $(\mathcal{Y}, \mathcal{D})$  be a family of extended pairs over  $S$ . A family of (stable) relative ideal sheaves of type  $\Gamma$  on  $(\mathcal{Y}, \mathcal{D})$  is an ideal sheaf  $\mathcal{I}$  of a closed subscheme  $Z \subset \mathcal{Y}$  such that  $\mathcal{O}_Z$  is flat over  $S$ , and for every closed point  $s \in S$ , the restriction  $\mathcal{I}_s = \mathcal{I} \otimes_{\mathcal{O}_S} k_s$  is a (stable) relative ideal sheaf on the extended pair  $(\mathcal{Y}_s, \mathcal{D}_s)$  of type  $\Gamma$ .

Next we define the moduli groupoid  $\mathfrak{I}_{Y,D}^\Gamma$  of stable relative ideal sheaves of type  $\Gamma$ . An object in the category  $\mathfrak{I}_{Y,D}^\Gamma$  is a family of stable relative ideal sheaves of type  $\Gamma$  over an extended pairs  $(\mathcal{Y}, \mathcal{D})$  parameterized by  $S$ . Let  $\mathcal{I}_1$  and  $\mathcal{I}_2$  be two families over  $(\mathcal{Y}_1, \mathcal{D}_1)$  and  $(\mathcal{Y}_2, \mathcal{D}_2)$  respectively. Then an arrow  $r : \mathcal{I}_1 \rightarrow \mathcal{I}_2$  consists of an arrow  $u : (\mathcal{Y}_1, \mathcal{D}_1) \rightarrow (\mathcal{Y}_2, \mathcal{D}_2)$  in stack  $\mathfrak{Y}$ , and an isomorphism of ideal sheaves  $\mathcal{I}_1 \cong u^* \mathcal{I}_2$  on  $\mathcal{Y}_1$ . Finally, there is a natural functor  $F : \mathfrak{I}_{Y,D}^\Gamma \rightarrow \text{Sch}$  which sends a family of ideal sheaves  $\mathcal{I}$  to the scheme  $S$ . It is direct to verify that the category  $\mathfrak{I}_{Y,D}^\Gamma$  and the functor  $F$  form a groupoid. Similarly, one can show that this groupoid is in fact a stack.

**Theorem 3.7.** The stack  $\mathfrak{I}_{Y,D}^\Gamma$  is a separated and proper Deligne-Mumford stack.

*Proof.* The proof is parallel to  $\mathfrak{I}_{X/C}^\Gamma$ . □

Let  $\iota : D \rightarrow Y[n]_0$  be the closed embedding of the distinguished divisor  $D[n]_0$ . It induces a homomorphism  $\iota^* : H^*(Y[n]_0) \rightarrow H^*(D)$ . Let  $c(\mathcal{I}) \in H^*(Y[n]_0)$  be the Chern class of a relative ideal sheaf  $\mathcal{I}$ . Then  $\iota^*(c)$  is the Chern class of  $\mathcal{I}_D$ . Let  $\Gamma$  be the fixed class in  $H_*(Y, \mathbf{Z})$ . Let  $\Gamma^\vee$  be the Poincare dual of  $\Gamma$ . We denote  $\Gamma_D$  the Poincare dual of  $\iota^*(\Gamma^\vee)$  in  $H_*(D, \mathbf{Z})$ . In fact,  $\iota_*(\Gamma_D)$  is the intersection of  $\Gamma$  and  $D$  in  $Y$ .

Let  $\mathcal{I}$  be a family of relative ideal sheaf of a closed subscheme  $Z$  on  $(\mathcal{Y}, \mathcal{D})$  parameterized by  $S$ . Then  $Z$  is flat over  $S$ . By the definition of relativeness, we know that the restriction of  $\mathcal{I}$  to  $\mathcal{D}$  is an ideal sheaf on  $\mathcal{D}$ .

**Proposition 3.8.** The assignment  $\mathcal{I} \mapsto \mathcal{I} \otimes_{\mathcal{O}_Y} \mathcal{O}_\mathcal{D}$  gives a natural morphism  $\mathfrak{I}_{Y,D}^\Gamma \rightarrow \mathbf{Hilb}^{\Gamma_D}(D)$ .

*Proof.* We show that this assignment is functorial. Let  $S$  be a scheme and  $\mathcal{I}$  be an object in  $\mathfrak{I}_{Y,D}^\Gamma(S)$ . Then  $\mathcal{I}$  is a family of stable relative ideal sheaves on an extended pair  $(\mathcal{Y}, \mathcal{D})$  parameterized by  $S$ . Since  $\mathcal{I}$  is perfect, the restriction  $\mathcal{I}_\mathcal{D} = \mathcal{I} \otimes_{\mathcal{O}_Y} \mathcal{O}_\mathcal{D}$  is an ideal sheaf on  $\mathcal{D}$ . By the argument in Lemma 2.29, we know that  $\mathcal{I}_\mathcal{D}$  is a family of ideal sheaves on  $\mathcal{D}$ , that is, the subscheme defined by  $\mathcal{I}_\mathcal{D}$  is flat over  $S$ . Note that  $\mathcal{D}$  is a constant family of  $D$  over  $S$ , we get a family of ideal sheaves on  $D$  parameterized by  $S$ . □

Let  $\xi_i = (\mathcal{Y}_i, \mathcal{D}_i)$  be families of extended pairs parameterized by  $S$  in  $\mathfrak{Y}_i$  respectively. Suppose  $\mathcal{D}_1 \cong \mathcal{D}_2$  as constant families over  $S$ . Let  $\mathcal{X}$  be the image of  $\xi_1 \times \xi_2$  under the morphism  $G : \mathfrak{Y}_1 \times_{\mathfrak{D}} \mathfrak{Y}_2 \rightarrow \mathfrak{X}_0$ . Then  $\mathcal{X}$  is a family of semistable models of  $X_0$  over  $S$ . Let  $\mathcal{I}_1$  and  $\mathcal{I}_2$  be families of stable relative ideal sheaves on  $(\mathcal{Y}_i, \mathcal{D}_i)$ . Suppose the restrictions of  $\mathcal{I}_1$  and  $\mathcal{I}_2$  to  $\mathcal{D}$  are isomorphic families of ideal sheaves, then by Lemma 2.13, there is a unique stable perfect ideal sheaf  $\mathcal{I}$  on  $\mathcal{X}$  such that the restriction of  $\mathcal{I}$  to  $(\mathcal{Y}_i, \mathcal{D}_i)$  is isomorphic to  $\mathcal{I}_i$ . This way, we obtain a morphism

$$\Phi : \mathfrak{I}_{Y_1, D_1}^{\Gamma_1} \times_{\mathbf{Hilb}^{\Gamma_D}(D)} \mathfrak{I}_{Y_2, D_2}^{\Gamma_2} \longrightarrow \mathfrak{I}_{X/C}^\Gamma(\mathbf{0}).$$

**Proposition 3.9.** *The morphism  $\Phi$  is a closed embedding of Deligne-Mumford stacks.*

*Proof.* It is direct to verify that  $\Phi$  is an isomorphism of  $\mathfrak{J}_{Y_1, D_1}^{\Gamma_1} \times_{\mathbf{Hilb}^{\Gamma_D}(D)} \mathfrak{J}_{Y_2, D_2}^{\Gamma_2}$  with the image stack which is a closed substack of  $\mathfrak{J}_{X/C}^{\Gamma}(\mathbf{0})$ .  $\square$

#### 4. PROOF OF LEMMA 2.29

Let  $R = k[[t]]$  be a complete discrete valuation ring. Let  $A = k[[t, y, x_i]]$  and  $A' = k[[y_1, y_2, x_i]]$  be the rings of formal power series. Define a homomorphism  $R \rightarrow A'$  by  $t \mapsto y_1 y_2$  and a natural injection  $R \rightarrow A$ . Then  $A$  and  $A'$  are flat  $R$ -algebras. Geometrically, it says that  $X = \text{Spec } A'$  is a flat family of affine formal schemes over  $T = \text{Spec } R$  and  $Y = \text{Spec } A$  is a constant family over  $T$ . We use  $X_0$  to denote the fiber of  $X \rightarrow T$  over the closed point of  $T$  and  $X_1$  the fiber over the generic point of  $T$ . Let  $k$  be the residue field of  $R$  and  $K$  be the field of fractions of  $R$ . Then  $X_0 = \text{Spec } k[[y_1, y_2, x_i]]/(y_1 y_2)$ . We let  $D$  be a closed subscheme of  $X$  defined by the ideal  $(y_1, y_2) \subset A$ . Likewise, we use  $Y_0$  and  $Y_1$  to denote the special and generic fiber of  $Y \rightarrow T$ . Let  $D$  be the closed subscheme of  $Y$  define by the ideal  $(y) \subset A$ . We also let  $D_0$  and  $D_1$  the corresponding closed subschemes of  $Y_0$  and  $Y_1$  respectively.

Let  $I \subset A$  or  $A'$  be an ideal. We can similarly denote  $I_0 = I \otimes_R k$  and  $I_1 = I \otimes_R K$ . Now we assume that  $I$  satisfies the following two conditions:

- (1)  $A/I$  or  $A'/I$  is flat over  $R$ .
- (2) The ideal  $I_1$  is relative with respect to  $D_1$  if  $I \subset A'$ .

In general, if  $I \subset A'$ ,  $I_0$  is not perfect as an ideal sheaf on  $X_0$ . If  $I \subset A$ ,  $I_0$  is not relative to  $D_0$ . We will show that, probably after a base change  $q : T' \rightarrow T$ ,  $X_1 \times_T T'$  or  $Y_1 \times_T T'$  can be extended to new families  $X' \rightarrow T'$  or  $Y' \rightarrow T'$ , such that the special fibre  $X'_0$  is a semistable model of  $X_0$  or  $(Y'_0, D'_0)$  is an extended pair of  $(Y_0, D_0)$ , and the family of ideal sheaves  $q^* I_1$  over the generic point  $\eta' \in T'$  can be extended to a family of ideal sheaves  $I'$  over  $X'/T'$  or  $Y'/T'$ , so that  $I'_0$  is perfect or relative according to  $I \subset A'$  or  $I \subset A$ , and  $I'_0$  has no trivial component.

Indeed, Lemma 2.29 gives a constructive method to find such an extension. Now we work out the local coordinate expression of the group action of  $G[n]$ .

Let  $S$  be a smooth curve with a distinguished closed point  $\eta_0$  and  $q : S \rightarrow C$  be a morphism such that  $q(\eta_0) = \mathbf{0}$ . We may also take  $S = \text{Spec } R$  to be a discrete valuation ring  $R$ . Let  $\mathcal{X}$  be an effective degeneration over  $S$  associated to a morphism  $f : S \rightarrow C[n]$ , such that  $f(\eta_0) = \mathbf{0} \in C[n]$ . Then it is clear that  $\mathcal{X}_0 = X[n]_0$ . Let  $\iota_k : C[n] \rightarrow C[n+1]$  be a standard embedding associated to  $\{1, \dots, \hat{k}, \dots, n+2\}$ . Let  $\rho : S - \{\eta_0\} \rightarrow G[n+1]$  be a morphism such that  $f^\rho = (\iota_k \circ f)^\rho : S - \{\eta_0\} \rightarrow C[n+1]$  can be extended to  $S$  and  $f^\rho(\eta_0) = \mathbf{0} \in C[n+1]$ . The morphism  $f^\rho : S \rightarrow C[n+1]$  induces an effective degeneration  $\mathcal{X}'$  over  $S$  such that  $\mathcal{X}' \times_S S^* \cong \mathcal{X} \times_S S^*$ , where  $S^* = S - \{\eta_0\}$ .

Let  $s$  be a local uniformizer of  $S$  at  $\eta_0$ . To illustrate the idea, we now assume  $n = 0$  and let the morphism  $f : S \rightarrow C$  be given by

$$t_1 = s^{\alpha_1},$$

and  $\rho : S^* \rightarrow G[1]$  is defined as

$$\sigma_1 = s^{\beta_1}.$$



Let  $p \in D$ . Then there are local coordinate functions  $y_1, y_2$  around  $p$  in  $X$  satisfying  $y_1 y_2 = t_1$ . Under the standard embedding,  $p$  can be regarded as a point in  $X[1]$ , the coordinates around  $p$  in  $X[1]$  are  $t_1, 1, y_1, y_2$ , satisfying

$$y_1 y_2 = t_1.$$

Apply the group action  $\rho$ , a point  $Q$  with coordinate  $(t_1, 1, y_1, y_2)$  was sent to the following point

$$Q^\rho = (\sigma_1 t_1, \sigma_1^{-1} \cdot 1, y_1, y_2).$$

Since we assume the morphism  $f^\rho : S^* \rightarrow C[1]$  can be extended to  $S$  and  $f^\rho(\eta_0) = \mathbf{0}$ ,  $Q^\rho \rightarrow \mathbf{0}$  when  $s \rightarrow 0$ . Recall the desingularization in section 1, we have local equations

$$u_1 y_1 = u_2 t_1, \quad u_2 y_2 = u_1 t_2,$$

where  $u_1, u_2$  are homogeneous coordinates on  $\Delta_1 \subset X[1]_0$ . Since the restriction of  $X[1]$  to  $t_2$ -axes is a smooth of  $X[1]_0$  along divisor  $D_2$ , we may find the unique point  $p \in D_1$  corresponding to the given point  $p \in D$ . The homogeneous coordinate of  $p$  is  $(u_1, u_2) = (0, 1)$ . Substitute the coordinate of  $Q^\rho$  into the two defining equations and set  $u_2 = 1$ , we obtain

$$\begin{aligned} u_1 y_1 = t_1 &= \sigma_1 t_1 = s^{\beta_1} s^{\alpha_1} = s^{\alpha_1 + \beta_1}, \\ y_2 = u_1 t_2 &= u_1 \sigma_1^{-1} \cdot 1 = u_1 s^{-\beta_1}. \end{aligned}$$

After a base change  $\tilde{s} = s^{\alpha_1 + \beta_1}$ , we have

$$(1) \quad y_2 = u_1 s^{-\beta_1} = u_1 \tilde{s}^{\frac{-\beta_1}{\alpha_1 + \beta_1}}.$$

Followed by this fact, we will forget the standard models and group actions in Lemma 2.29. The essence of the group action is the substitution

$$y = t^a w.$$

for positive rational number  $a$ . Notice that this substitution is a combination of the process of blowup and base change, since the local expression for blowup is  $y = tw$ , and for base change is  $t' = t^n$ .

In what follows, a substitution  $y = t^a w$  for positive rational number  $a = \frac{p}{q}$  always means applying base change  $t \mapsto t^q$  first and then blow up  $p$  times. We can actually apply this substitution formally and forget the geometric interpretation.

In the remainder of this section, we prove Lemma 2.29 for the case of ring  $A$ . The proof of the case of ring  $A'$  can be reduced to similar situation as ring  $A$  as follows. Since  $X = \text{Spec } A'$  is a family over  $T$  corresponding to the homomorphism  $k[[t]] \rightarrow A' = k[[y_1, y_2, x_i]]$  given by  $t \mapsto y_1 y_2$ , we can actually regard  $X$  as a family over  $\text{Spec } k[[y_1]]$  using the injection  $k[[y_1]] \rightarrow A'$ . This family is generically flat. Now we think of  $y_1$  as the parameter  $t$  in the case of ring  $A$ . By (1), the substitution becomes

$$y_2 = u_1 \tilde{s}^{\frac{-\beta_1}{\alpha_1 + \beta_1}} = u_1^{\frac{\alpha_1}{\alpha_1 + \beta_1}} y_1^{\frac{-\beta_1}{\alpha_1 + \beta_1}}.$$

Put it simply, it is a substitution of the form  $y = t^a w^b$ . Therefore, the results in the remainder for the substitution  $y = t^a w$  can be generalized to this case directly.

Now we focus on the substitution  $y = t^a w$ . To begin with, we introduce some notation.

Let  $B = k[[t, w, x_i]]$  and let  $\bar{B} \subset B$  be a subring of  $B$  such that for any element  $f(t, w, x_i) \in \bar{B}$ ,  $f(0, w, x_i)$  is a polynomial of  $w$  with coefficients in  $k[[x_i]]$ . It is clear that  $k[[t, x_i]][w] \subset \bar{B}$ .

**Definition 4.1.** Let  $A = k[[t, y, x_i]]$  and  $I \subset A$  be an ideal. Let  $f : A \rightarrow \bar{B}$  be a homomorphism defined by  $f(y) = t^a w$ . Then we denote by  $I(a)$  the ideal in  $\bar{B}$  generated by  $f(I)$ .

It is easy to see that for any  $f(t, w, x_i) \in \bar{B}$ , there is an integer  $n$  and an element  $g(t, y, x_i) \in A$ , such that  $t^n f(t, w, x_i) = g(t, t^a w, x_i)$ . In what follows, we let  $B_0 = k[[w, x_i]]$  and  $\bar{B}_0 = \bar{B} \cap B_0 = k[[x_i]][w]$ . For relativeness, it is not important to distinguish  $\bar{B}$  and  $B$ , that is, we can define  $I(a)$  to be an ideal in  $B$ . However, when considering trivial component of an ideal, we have to work on  $\bar{B}$  rather than  $B$ .

Let  $R = k[[t]]$  and  $\iota_B : R \rightarrow B$  be the canonical injective homomorphism. For any ideal  $I \subset R$ , we use  $I_B$  or  $I[[w, x_i]]$  to denote the ideal in  $B$  generated by the  $\iota_B(I)$ . A standard fact in commutative algebra says that for any nonzero element  $b$  in  $B$ , there is a unique nonnegative integer  $n$  and a unique nonzero element  $b_0$  in  $k[[w, x_i]]$ , such that  $b - b_0 t^n$  is contained in  $\mathfrak{m}_B^{n+1}$ . In the sequel,  $b_0$  is called the leading coefficient of  $b$ .

**Definition 4.2.** Let  $J \subset B$  be an ideal. The initial ideal of  $J$ , denoted by  $L(J)$ , is defined to be the set of leading coefficients of elements in  $J$ .

**Definition 4.3.** Given  $a > 0$  and an ideal  $I(a) \subset \bar{B}$ , we denote  $\widetilde{I(a)}$  to be the ideal consists of elements  $b$  in  $\bar{B}$  such that  $t^n b \in I(a)$  for some integer  $n \geq 0$ .

It is obvious from the definition that  $I(a) \subset \widetilde{I(a)}$ .

**Lemma 4.4.** The  $R$ -algebra  $B/\widetilde{I(a)}$  is flat over  $R$ .

Let  $\widetilde{I(a)}_0 = \widetilde{I(a)} \otimes_R k$  be the fiber of  $\widetilde{I(a)}$  over the closed point  $\eta_0 \in T$ . Then  $\widetilde{I(a)}_0$  is exactly the initial ideal of  $I(a)$ . Recall that  $A = k[[t, y, x_i]]$  and  $Y = \text{Spec } A$ , and  $D \subset Y$  is a divisor defined by  $y = 0$ ,  $D = \text{Spec } k[[t, x_i]]$  and there is a surjective homomorphism of rings  $\pi : A \rightarrow k[[t, x_i]]$  given by  $\pi(y) = 0$  corresponding to the closed embedding  $\iota : D \rightarrow Y$ . Let  $I \subset A$  be the ideal of a closed subscheme  $Z \subset Y$ . Since  $\pi$  is surjective, the image  $\pi(I)$  is an ideal of  $k[[t, x_i]]$  which is denoted by  $I^D$ . Then  $I^D$  is the ideal of the closed subscheme  $\iota^{-1}(Z) = Z \cap D \subset D$ . Let  $I_{t \rightarrow 0}^D \subset k[[x_i]]$  be the initial ideal of  $I^D$ . Let  $I_0^D$  be the fiber of  $I^D$  at the closed point  $\eta_0 \in T$ . We always have  $I_0^D \subset I_{t \rightarrow 0}^D$ .

Let  $I \subset A$  be an ideal. Since there is a surjective homomorphism  $A \rightarrow k[[y, x_i]]$  which sends  $t$  to 0, the fiber  $I_0 = I \otimes_R k$  is exactly the image of  $I$  in  $k[[y, x_i]]$  via the previous homomorphism.

**Lemma 4.5.**  $I_0$  is relative to  $D_0$  if and only if  $I_0^D = I_{t \rightarrow 0}^D$ .

*Proof.* We first prove the necessity, that is, if  $I_0$  is relative to  $D_0$ , then  $I_0^D = I_{t \rightarrow 0}^D$ . Because there is always an inclusion  $I_0^D \subset I_{t \rightarrow 0}^D$ , it suffices to show that  $I_0^D \supset I_{t \rightarrow 0}^D$ . Let  $f(x_i)$  be an element in  $I_{t \rightarrow 0}^D$ . By the definition of the ideal  $I_{t \rightarrow 0}^D$ , there exists at least one element  $h$  in the ideal  $I^D$  such that  $f(x_i)$  is the leading coefficient of  $h$ , that is

$$h - f(x_i)t^l \in \mathfrak{m}^{l+1}[[x_i]]$$

with  $l \geq 0$ . Let  $L_f$  be the subset of  $I^D$  consists of elements with leading coefficient  $f(x_i)$ . Let  $l_0$  be the least power of  $t$  in leading terms among all elements in  $L_f$ . We will show that  $l_0 = 0$ . If this has been proved, then there is an element in  $L_f$  of the form  $f(x_i) + g$  with  $g \in \mathfrak{m}[[x_i]]$ . It implies directly that  $f(x_i) \in I_0^D$ .

Now we show that  $l_0 = 0$ . Suppose not, then  $l_0 > 0$ . Let  $h$  be an element in  $L_f$  which has the lowest power  $l_0$  of  $t$  in the leading term. Then  $h = f(x_i)t^{l_0} + g_0$  with  $g_0 \in \mathfrak{m}^{l_0+1}[[x_i]]$ . We know  $h \in \mathfrak{m}[[x_i]]$  by  $l_0 > 0$ . Since there is a surjective homomorphism  $\pi : I \rightarrow I^D$ , we let

$$\alpha = h + y(g_1(y, x_i) + g_2), \quad g_2 \in \mathfrak{m}[[y, x_i]]$$

be an element in the inverse image of  $h$  in  $I$ . Now consider the image of  $\alpha$  via the homomorphism  $\psi : I \rightarrow I_0$ ,

$$\psi(\alpha) = \psi(h) + \psi(yg_1(y, x_i)) + \psi(yg_2).$$

Since  $\psi(\mathfrak{m}[[y, x_i]]) = 0$  and  $h \in \mathfrak{m}[[x_i]]$ ,  $yg_2 \in \mathfrak{m}[[y, x_i]]$ , we get  $\psi(\alpha) = \psi(yg_1(y, x_i)) = yg_1(y, x_i)$ . It implies that  $yg_1(x_i, y) \in I_0$ . By the relativeness of  $I_0$  with respect to  $D_0$  and Proposition 2.14,  $g_1(x_i, y)$  is also contained in  $I_0$ . So we can find an element in  $I$  of the form

$$\beta = g_1(x_i, y) + u, \quad u \in \mathfrak{m}[[y, x_i]].$$

Now consider the element  $\alpha - y\beta$  in  $I$ , notice that

$$\alpha - y\beta = h + y(g_2 - u) \in \mathfrak{m}[[y, x_i]],$$

we let  $g_0 = tg_3$  and  $g_2 - u = tg_4$  for some  $g_3 \in \mathfrak{m}^{l_0}[[x_i]]$  and  $g_4 \in A$ . Then

$$\alpha - y\beta = t(f(x_i)t^{l_0-1} + g_3 + yg_4).$$

Since  $A/I$  is flat over  $R$ ,  $t(f(x_i)t^{l_0-1} + g_3 + yg_4) \in I$  implies that  $f(x_i)t^{l_0-1} + g_3 + yg_4$  is also contained in  $I$ . The image of this element in  $I^D$ , under the homomorphism  $\pi$ , is  $f(x_i)t^{l_0-1} + g_3$ . It is an element with leading coefficient  $f(x_i)$ . However, the power of  $t$  in the leading term is  $l_0 - 1 \geq 0$ , which contradicts to the assumption that  $l_0$  is the least power. Therefore we have  $l_0 = 0$ .

Next we prove the sufficiency, that is, show that  $I_0$  is relative to  $D_0$  under the assumption  $I_0^D = I_{t \rightarrow 0}^D$ . By proposition 2.14, it suffices to show that  $yf(x_i, y) \in I_0$  implies that  $f(x_i, y) \in I_0$ . Let  $yf(x_i, y)$  be an element in  $I_0$ . Then there is an element

$$\alpha = yf(x_i, y) + t(g_1(x_i) + yg_2(y, x_i))$$

in  $I$ . It implies that  $tg_1(x_i)$  is contained in  $I^D$ . Since  $I_0^D = I_{t \rightarrow 0}^D$ , we get that  $k[[t, x_i]]/I^D$  is flat over  $R$ . Hence  $g_1(x_i)$  is contained in  $I^D$ . Therefore there exists  $h \in I$  such that

$$\beta = g_1(x_i) + yh$$

is contained in  $I$ . Consider  $\alpha - t\beta$  in  $I$ , because

$$\begin{aligned} \alpha - t\beta &= yf(x_i, y) + tyg_2 - tyh \\ &= y(f(x_i, y) + tg_2 - th), \end{aligned}$$

and the generic relativeness of  $I$ ,  $f(x_i, y) + tg_2 - th$  is also contained in  $I$ . It implies that  $f(x_i, y) \in I_0$ .

□

**Proposition 4.6.**  $\widetilde{I(a)_0}^D \subset I_{t \rightarrow 0}^D$  for any  $a > 0$ .

*Proof.* Let  $f_1(x_i), \dots, f_n(x_i)$  be a set of generators of the ideal  $\widetilde{I(a)_0}^D$ . Then there exist  $\tilde{f}_1(w, x_i), \dots, \tilde{f}_n(w, x_i)$  in  $\widetilde{I(a)_0}$ , such that

$$\tilde{f}_k(w, x_i) = f_k(x_i) + wg_k(w, x_i)$$

for  $k = 1, \dots, n$ . Since  $\widetilde{I(a)}_0$  is the initial ideal of  $I(a)$ , we can find elements  $g_1(t, w, x_i), \dots, g_n(t, w, x_i) \in I(a)$ , such that

$$g_k(t, w, x_i) = t^{n_k} \tilde{f}_k(w, x_i) + t^{n_k + \mu_k} h_k(t, w, x_i)$$

for some  $n_k \geq 0$  and  $\mu_k > 0$ .

Now let  $w = 0$  in  $g_k(t, w, x_i)$ . We know that the elements

$$g_k(t, 0, x_i) = t^{n_k} \tilde{f}_k(0, x_i) + t^{n_k + \mu_k} h_k(t, 0, x_i)$$

are contained in  $I(a)^D = I^D$ . The leading coefficient of  $g_k(t, 0, x_i)$  is  $f_k(0, x_i) = f_k(x_i)$ , it implies that  $f_k(x_i)$  is contained in  $I_{t \rightarrow 0}^D$ . Since  $\widetilde{I(a)}_0^D = (f_1(x_i), \dots, f_n(x_i))$ , we obtain  $\widetilde{I(a)}_0^D \subset I_{t \rightarrow 0}^D$ .  $\square$

Now let  $0 < a < b$ , we obtain a sequence

$$I_0^D \subset \widetilde{I(a)}_0^D \subset \widetilde{I(b)}_0^D \subset I_{t \rightarrow 0}^D.$$

By Lemma 4.5,  $\widetilde{I(a)}_0$  is relative to  $D_0$  if and only if  $\widetilde{I(a)}_0^D = I_{t \rightarrow 0}^D$ . It follows that if  $\widetilde{I(a)}_0$  is relative to  $D_0$ , then  $\widetilde{I(b)}_0$  is also relative to  $D_0$  for any  $b > a$ .

Let  $q : \text{Spec } \bar{B}_0 \rightarrow \text{Spec } k[[x_i]]$  be the morphism corresponding to the injection  $k[[x_i]] \rightarrow \bar{B}_0$ . For an ideal  $J \subset k[[x_i]]$ , we define the pullback  $q^*J$  as the ideal  $J\bar{B}_0 \subset \bar{B}_0$ .

**Lemma 4.7.** *If  $q^*(I_{t \rightarrow 0}^D) \subset \widetilde{I(a)}_0$ , then  $q^*(I_{t \rightarrow 0}^D) = \widetilde{I(a)}_0$ .*

*Proof.* It suffices to show that  $\widetilde{I(a)}_0 \subset q^*(I_{t \rightarrow 0}^D)$ .

Let  $f(w, x_i) = f_0(x_i) + wf_1(w, x_i)$  be an element in  $\widetilde{I(a)}_0$ . Then there exists an element  $g(t, w, x_i) \in I(a)$ , such that

$$\begin{aligned} g(t, w, x_i) &= t^n f(w, x_i) + t^{n+\mu} g_1(t, w, x_i) \\ &= t^n (f_0(x_i) + wf_1(w, x_i)) + t^{n+\mu} g_1(t, w, x_i). \end{aligned}$$

It implies that

$$g(t, 0, x_i) = t^n f_0(x_i) + t^{n+\mu} g_1(t, 0, x_i)$$

is contained in  $I(a)^D$ . Since  $I_{t \rightarrow 0}^D$  is the initial ideal of  $I^D$  and  $I(a)^D = I^D$  for any  $a > 0$ ,  $I_{t \rightarrow 0}^D$  is also the initial ideal of  $I(a)^D$ . Therefore,  $g(t, 0, x_i) \in I(a)^D$  implies that  $f_0(x_i) \in I_{t \rightarrow 0}^D$ .

Since  $q^*(I_{t \rightarrow 0}^D) \subset \widetilde{I(a)}_0$  and  $f_0(x_i) \in q^*(I_{t \rightarrow 0}^D)$ , we get  $f_0(x_i) \in \widetilde{I(a)}_0$ . Hence  $wf_1(w, x_i) = f(w, x_i) - f_0(x_i)$  is also contained in  $\widetilde{I(a)}_0$ . On the other hand, since  $q^*(I_{t \rightarrow 0}^D) \subset \widetilde{I(a)}_0$ , we obtain  $I_{t \rightarrow 0}^D \subset \widetilde{I(a)}_0^D$ . It implies that  $\widetilde{I(a)}_0$  is relative to  $D_0$ . Therefore,  $wf_1(w, x_i) \in \widetilde{I(a)}_0$  implies that  $f_1(w, x_i) \in \widetilde{I(a)}_0$ . Continue the process, we get  $\widetilde{I(a)}_0 \subset q^*(I_{t \rightarrow 0}^D)$ .  $\square$

**Lemma 4.8.** *Let  $I_{t \rightarrow 0}^D = (g_1(x_i), \dots, g_n(x_i))$ . Then  $q^*(I_{t \rightarrow 0}^D) = \widetilde{I(a)}_0$  if and only if there exist  $f_1(t, y, x_i), \dots, f_n(t, y, x_i)$  in  $I$ , such that*

$$f_k(t, y, x_i) = t^{a_k} g_k(x_i) + t^{a_k + \mu_k} \tilde{g}_k(t, x_i) + y \tilde{h}_k(t, y, x_i),$$

*and the leading coefficient of  $f_k(t, t^a w, x_i)$  is exactly  $g_k(x_i)$ .*

*Proof.* We first show the “only if” part. Since  $q^*(I_{t \rightarrow 0}^D) = (g_i(x_i), \dots, g_n(x_i))$  and  $q^*(I_{t \rightarrow 0}^D) = \widetilde{I(a)}_0$ , every  $g_k(x_i)$  is contained in  $\widetilde{I(a)}_0$ . It implies that there are elements  $\tilde{g}_k(t, w, x_i) \in I(a)$  whose leading coefficients are  $g_k(x_i)$ . Now for  $\tilde{g}_k(t, w, x_i) \in I(a)$ , there exists an integer  $m$  such that  $t^m \tilde{g}_k(t, w, x_i) = f(t, t^a w, x_i)$  for some  $f(t, y, x_i) \in I$ . Let

$$f_k(t, y, x_i) = t^{a_k} h_k(x_i) + t^{a_k + \mu_k} \tilde{g}_k(t, x_i) + y \tilde{h}_k(t, y, x_i).$$

By the comparison of leading terms, we have  $g_k(x_i) = h_k(x_i)$ . Obviously,  $f_k(t, y, x_i) \in I$  satisfy the required property.

Next, we prove “if” part. Let  $I_{t \rightarrow 0}^D = (g_1(x_i), \dots, g_n(x_i))$ . By hypothesis, for each  $g_k(x_i)$ , there is an element  $f_k(t, y, x_i) \in I$

$$f_k(t, y, x_i) = t^{a_k} g_k(x_i) + t^{a_k + \mu_k} \tilde{g}_k(t, x_i) + y \tilde{h}_k(t, y, x_i),$$

such that the leading coefficient of  $f_k(t, t^a w, x_i)$  is  $g_k(x_i)$ . Since  $f_k(t, t^a w, x_i)$  is contained in  $I(a)$ , its leading coefficient  $g_k(x_i)$  is contained in  $\widetilde{I(a)}_0$ . Hence  $q^*(I_{t \rightarrow 0}^D) \subset \widetilde{I(a)}_0$  and the result follows from the previous lemma.  $\square$

**Corollary 4.9.**  $\widetilde{I(a)}_0$  is relative to  $D_0$  when  $a$  is sufficiently large.

*Proof.* By Lemma 4.5 and Lemma 4.6, it suffices to show that  $\widetilde{I(a)}_0^D \supset I_{t \rightarrow 0}^D$ .

The argument is same as the previous lemma. Let  $I_{t \rightarrow 0}^D = (g_1(x_i), \dots, g_n(x_i))$ . Then for each  $g_k(x_i)$ , there is an element  $f_k(t, y, x_i) \in I$

$$f_k(t, y, x_i) = t^{a_k} g_k(x_i) + t^{a_k + \mu_k} \tilde{g}_k(t, x_i) + y \tilde{h}_k(t, y, x_i),$$

If we choose  $a > \max\{a_1, \dots, a_n\}$ , then the leading coefficient of  $f_k(t, t^a w, x_i) \in I(a)$  is  $g_k(x_i)$ . Hence  $g_k(x_i)$  is contained in  $\widetilde{I(a)}_0$ . Note that  $g_k(x_i)$  does not depend on  $t$ , we get  $g_k(x_i) \in \widetilde{I(a)}_0^D$ .  $\square$

**Lemma 4.10.** If  $\widetilde{I(a)}_0$  is relative to  $D_0$ , then  $\widetilde{I(b)}_0 = q^*(I_{t \rightarrow 0}^D)$  for any  $b > a$ .

*Proof.* By Lemma 4.7, it suffices to show that  $q^*(I_{t \rightarrow 0}^D) \subset \widetilde{I(b)}_0$ . Since  $\widetilde{I(a)}_0$  is relative to  $D_0$ ,  $\widetilde{I(a)}_0^D = I_{t \rightarrow 0}^D$ . Let  $I_{t \rightarrow 0}^D = (g_1(x_i), \dots, g_n(x_i))$ . Since  $g_k(x_i) \in \widetilde{I(a)}_0^D$ , there exist  $\tilde{f}(t, w, x_i) \in I(a)$  such that

$$\tilde{f}(t, w, x_i) = t^{b_k} (g(x_i) + w \tilde{g}_k(w, x_i)) + t^{b_k + \mu_k} \tilde{h}(t, w, x_i).$$

Since  $\tilde{f}(t, w, x_i) \in I(a)$ , there is an integer  $m$  and an element  $f(t, y, x_i) \in I$ , such that  $t^m \tilde{f}(t, w, x_i) = f(t, t^a w, x_i)$ . Now for any  $b > a$ , make a substitution  $w \mapsto t^{b-a} w$  in  $\tilde{f}(t, w, x_i)$ , we get

$$\begin{aligned} f(t, t^b w, x_i) &= \tilde{f}(t, t^{b-a} w, x_i) \\ &= t^{b_k} (g(x_i) + t^{b-a} w \tilde{g}_k(t^{b-a} w, x_i)) + t^{b_k + \mu_k} \tilde{h}(t, t^{b-a} w, x_i), \end{aligned}$$

whose leading coefficient is  $g_k(x_i)$ . By Lemma 4.8,  $q^*(I_{t \rightarrow 0}^D) = \widetilde{I(b)}_0$ .  $\square$

Now we define the set

$$S = \{a_0 \in \mathbf{R} \mid \widetilde{I(a)}_0 = q^*(I_{t \rightarrow 0}^D) \text{ for } a > a_0\}.$$

By Corollary 4.9 and Lemma 4.10, the set  $S$  is not empty. Let  $a_* = \inf S$ . We have

**Proposition 4.11.** (1)  $a_*$  is a positive rational number.

(2) The ideal  $\widetilde{I(a_*)}_0$  is relative to  $D_0$ .

(3)  $\widetilde{I(a_*)}_0 \neq q^*(I_{t \rightarrow 0}^D)$ .

*Proof.* (1) Let  $I_{t \rightarrow 0}^D = (g_1(x_i), \dots, g_n(x_i))$ . From the definition of  $a_*$ , we know that for any rational number  $b > a_*$ ,  $\widetilde{I(b)}_0 = q^*(I_{t \rightarrow 0}^D)$ . By Lemma 4.8, there are elements  $f_k(t, y, x_i) \in I$  such that

$$f_k(t, y, x_i) = t^{a_k} g_k(x_i) + t^{a_k + \mu_k} \tilde{g}_k(t, x_i) + y \tilde{h}_k(t, y, x_i),$$

and the leading coefficient of  $f_k(t, t^b w, x_i)$  in  $I(b)$  is  $g_k(x_i)$ . Now we write  $\tilde{h}_k(t, y, x_i)$  as  $\sum c_{lm}(x_i) t^l y^m$ , then

$$\tilde{h}_k(t, t^b w, x_i) = \sum c_{lm}(x_i) t^l (t^b w)^m = \sum c_{lm}(x_i) t^{l+mb} w^m.$$

Therefore, when  $y = t^b w$ , the leading term of  $y \tilde{h}_k(t, y, x_i)$  has the form

$$(t^b w) c_{lm}(x_i) t^{l+mb} w^m = c_{lm}(x_i) t^{l+(m+1)b} w^{m+1},$$

in which the power of  $t$  is  $l + (m + 1)b$  and  $l, m$  are both nonnegative rational numbers.

Since for  $b > a_*$ , the leading coefficient of  $f_k(t, t^b w, x_i)$  is  $g_k(x_i)$ , we get  $l + (m + 1)b > a_k$ . By the continuity of  $l + (m + 1)b$  as a function of  $b$ , we obtain  $l + (m + 1)a_* \geq a_k$ . Suppose  $a_*$  is not rational, the equality does not hold since  $a_k$  is rational. We can find rational number  $\tilde{a} < a_*$  such that  $l + (m + 1)\tilde{a} > a_k$  for all  $k$ . It implies that the leading coefficient of  $f_k(t, t^{\tilde{a}} w, x_i)$  is  $g_k(x_i)$ . By Lemma 4.8,  $\widetilde{I(\tilde{a})}_0 = q^*(I_{t \rightarrow 0}^D)$ . From Lemma 4.10, we get  $\widetilde{I(b)}_0 = q^*(I_{t \rightarrow 0}^D)$  for all  $b > \tilde{a}$ . Since  $\tilde{a} < a_*$ , it contradicts to the definition of  $a_*$ .

(2) It suffices to show that  $I_{t \rightarrow 0}^D \subset \widetilde{I(a_*)}_0^D$ . Let  $I_{t \rightarrow 0}^D = (g_1(x_i), \dots, g_n(x_i))$ . We will show that  $g_k(x_i) \in \widetilde{I(a_*)}_0^D$  for all  $k$ . Following the notation in part (1), we see that the leading coefficient of  $f_k(t, t^{a_*} w, x_i)$  is  $g_k(x_i)$  if  $l + (m + 1)a_* > a_k$ , and  $g_k(x_i) + c_{lm}(x_i) w^{m+1}$  if  $l + (m + 1)a_* = a_k$ . Notice that the leading coefficient  $g_k(x_i)$  or  $g_k(x_i) + c_{lm}(x_i) w^{m+1}$  is contained in  $\widetilde{I(a_*)}_0$ , we obtain  $g_k(x_i) \in \widetilde{I(a_*)}_0^D$  by setting  $w = 0$ .

(3) Suppose  $\widetilde{I(a_*)}_0 = q^*(I_{t \rightarrow 0}^D)$ , we will show that there exists a rational number  $\theta > 0$ , such that for any  $a > a_* - \theta$ ,  $\widetilde{I(a)}_0 = q^*(I_{t \rightarrow 0}^D)$ . It will imply that  $a_* - \theta$  is contained in  $S$  which violates the definition of  $a_* = \inf S$ .

We let  $I_{t \rightarrow 0}^D = (g_1(x_i), \dots, g_n(x_i))$ . Then by Lemma 4.8, there exist  $f_k(t, y, x_k) \in I$  such that

$$f_k(t, y, x_i) = t^{a_k} g_k(x_i) + t^{a_k + \mu_k} \tilde{g}_k(t, x_i) + y \tilde{h}_k(t, y, x_i),$$

and the leading coefficient of  $f_k(t, t^{a_*} w, x_i)$  is exactly  $g_k(x_i)$ .

If we choose a rational number  $\theta_k$  small enough, then by the continuity, for  $a > a_* - \theta_k$ , the leading coefficient of  $f_k(t, t^a w, x_i)$  is same as the leading coefficient of  $f_k(t, t^{a_*} w, x_i)$ . It implies that  $g_k(x_i)$  is contained in  $\widetilde{I(a)}_0$ . Now we let  $\theta =$

$\min\{\theta_1, \dots, \theta_n\}$ . Then for  $a > a_* - \theta$ ,  $(g_1(x_i), \dots, g_n(x_i))$  is contained in  $\widetilde{I(a)}_0$ . Now by Lemma 4.7, we obtain  $\widetilde{I(a)}_0 = q^*(I_{t \rightarrow 0}^D)$  for  $a > a_* - \theta$ .  $\square$

From Lemma 4.10, the rational number satisfying conditions (2) and (3) in this proposition is unique.

## REFERENCES

- [1] K. Behrend, and B. Fantechi, *The intrinsic normal cone*. Invent. Math. 128 (1997), no. 1, 45–88.
- [2] P. Deligne and D. Mumford, *The irreducibility of the space of curves of given genus*. Inst. Hautes tudes Sci. Publ. Math. No. 36(1969) 75–109.
- [3] S.K. Donaldson and R.P. Thomas, *Gauge theory in higher dimensions*. In: The geometric universe (Oxford, 1996), 31–47. Oxford Univ. Press, Oxford, 1998.
- [4] A. Grothendieck, *Techniques de descente et théorèmes d’existence en géométrie algébrique, IV*. Sem. Bourbaki, 221, 1960–1961.
- [5] M. Levine and R. Pandharipande, *Algebraic cobordism revisited*. Preprint, math.AG/0605196.
- [6] J. Li, *Virtual moduli cycles and Gromov-Witten invariants of algebraic varieties*. J. Amer. Math. Soc. 11 (1998), no. 1, 119–174.
- [7] J. Li, *Stable morphisms to singular schemes and stable relative morphisms*. J. Diff. Geom. 57(2001) 509–578.
- [8] J. Li, *Lecture notes on relative GW-invariants*. Intersection theory and moduli, 41–96 (electronic), ICTP Lect. Notes, XIX, Abdus Salam Int. Cent. Theoret. Phys., Trieste, 2004.
- [9] J. Li, *Zero dimensional Donaldson-Thomas invariants of threefolds*. math.AG/0604490.
- [10] D. Maulik, N. Nekrasov, A. Okounkov and R. Pandharipande, *Gromov-Witten theory and Donaldson-Thomas theory I*. math.AG/0312059.
- [11] D. Maulik, N. Nekrasov, A. Okounkov and R. Pandharipande, *Gromov-Witten theory and Donaldson-Thomas theory II*. math.AG/0406092.
- [12] R.P. Thomas, *A holomorphic Casson invariant for Calabi-Yau 3-folds, and bundles on K3 fibrations*. J. Differential Geom. 54 (2000), 367–438.
- [13] A. Vistoli, *Intersection theory on algebraic stacks and on their moduli spaces*. Invent. Math. 97(1989), no. 3, 613–670.

DEPARTMENT OF MATHEMATICS, STANFORD UNIVERSITY, STANFORD, CA 94305  
*E-mail address*: `bwu@math.stanford.edu`